

Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

## DELARO<sup>™</sup> 325 SC Fungicide

For suppression of white mold in soybean EPA Reg. No. 264-1055

## FIFRA 2(ee) Recommendation

## FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF CONNECTICUT, DELAWARE, ILLINOIS, INDIANA, IOWA, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA, NEBRASKA, NEW HAMPSHIRE, NEW JERSEY, NORTH DAKOTA, OHIO, PENNSYLVANIA, RHODE ISLAND, SOUTH DAKOTA, VERMONT, VIRGINIA, WEST VIRGINIA, AND WISCONSIN

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Disease Suppressed	Product Rate (fl oz/A)	Product Instructions
White mold <i>(Sclerotinia sclerotiorum)</i>	8.0 - 11.0	<ul> <li>Apply Delaro 325 SC Fungicide as a broadcast foliar spray at R1 (beginning bloom) when conditions are favorable for disease development before infection occurs.</li> <li>A sequential treatment of DELARO 325 SC Fungicide or Stratego YLD Fungicide may be made at R3 – R4 (beginning to full pod). DELARC 325 SC Fungicide may be applied by ground, air, or chemigation. Apply in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 3 gallons of spray solution per acre by aircraft spray equipment.</li> <li>Use of the higher rates are recommended when disease pressure is expected to be severe.</li> </ul>
Application Restrictions: Pre-Harvest Interval (PHI): <b>21 day(s)</b> Do not graze or feed soybean forage or hay. Do not apply more than 33.0 fl oz per acre p Do not exceed 0.53 lbs prothioconazole per Do not apply more than 3 applications per ye	er year. acre per year, or 0.33 lbs trifloxystrob	in per acre per year.

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577. For <u>PRODUCT</u> <u>USE</u> Information Call 1-866-99BAYER (1-866-992-2937).

> As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).