



**FIFRA 24 (C)
Special Local Need Label
For Use on Hops Only**

EPA Reg. No. 279-3241

EPA SLN. No. OH-140002

FIFRA Section 24 (c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OHIO.

This label is valid until September 3, 2019 or until otherwise amended, withdrawn, canceled, or suspended.

Aim EC Herbicide

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this Supplemental label and the main EPA-registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CROP/SITE:	TARGET PEST:	RATE:	METHOD OF APPLICATION:	FREQUENCY AND TIMING OF APPLICATION:
Hops	Apply Aim EC using shielded sprayers or hooded sprayers to control emerged and actively growing broadleaf weeds within or between the rows of the crop. Refer to weed control list below.	0.5 – 2.0 fl. oz. (0.008 – 0.032 pound active ingredient) per acre. Refer to weed control list below.	Apply Aim EC at 2.0 fl. oz. (0.03 pound active ingredient) per acre per application in a minimum of 20 gallons of spray solution by boom-type ground application equipment only to the basal portion of the hop plant (approximately the lower 1.5 feet) and to the sucker mat which extends from the base of the plant to approximately 1.5 to 2 feet into the row. An alternate row treatment program may be followed to avoid the removal of excessive photosynthetic capacity from the crown area. When treating alternating rows on different days, the equivalent maximum rate must not exceed 3.2 fl. oz. (0.015 pound active ingredient) of Aim EC per application per treated row area totaling 0.5 acres.	Do not apply more than 7.6 fl. oz. (0.12 pound active ingredient) per acre per season. Allow 14 days between treatments of Aim EC.

Use Restrictions:

- Do not apply Aim EC using air blast or air assisted sprayers.
- Do not apply within 7 days of harvest.
- Do not apply through any type of irrigation system.

This Condition of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Aim is a trademark of FMC Corporation.

©2014 All rights reserved.

Rev: 9-9-2014



**FMC Corporation
Agricultural Solutions
1735 Market Street
Philadelphia, PA 19103**

Weeds Controlled	Aim EC Use Rate fl. oz. (pound active ingredient) per acre
Lambsquarters, common (up to 3 inches tall)	0.5 fl. oz. (0.008 pound active ingredient) per acre
Morningglory, ivyleaf (up to 3 leaves)	
Morningglory, pitted (up to 3 leaves)	
Nightshade, Eastern black	
Pigweed, redroot	
Velvetleaf	
Waterhemp (up to 2 inches tall)	
Weeds Controlled	Aim EC Use Rate fl. oz. (pound active ingredient) per acre
All the weeds controlled at 0.5 fl. oz. (0.008 pound active) per acre plus the weeds listed below:	0.8 fl. oz. (0.013 pound active ingredient) per acre
Cheeseweed	
Filaree, redstem	
Flixweed	
Lambsquarters, common	
Morningglory, entireleaf	
Morningglory, ivyleaf	
Morningglory, pitted	
Morningglory, scarlet	
Nightshade, hairy	
Pennycress, field	
Pigweed, prostrate	
Pigweed, smooth	
Pigweed, tumble	
Purslane, common	
Sesbania, hemp	
Smartweed, PA (seedling)	
Spurge, prostrate	
Tansymustard	
Velvetleaf (24")	
Waterhemp, common & tall	
Weeds Controlled	Aim EC Use Rate fl. oz. (pound active ingredient) per acre
All the weeds controlled at 0.8 fl. oz. (0.013 pound active) per acre plus the weeds listed below:	1.0 fl. oz. (0.016 pound active ingredient) per acre
Amaranth, spiny	
Anoda, spurred	
Bedstraw, catchweed	
Buffalobur	
Carpetweed	
Cocklebur	
Copperleaf, hophornbeam	
Cotton, GMO Varieties	
Cotton, volunteer	
Eclipta	
Fiddleneck, coast	
Groundcherry, smooth (seedling)	
Groundcherry, Wright's	
Jimsonweed	
Kochia	
Lettuce, Prickly 2-3 leaf	
Nettle, burning	
Nightshade, American black	
Nightshade, black	
Rocket, London	
Shepherdspurse	
Speedwell, Virginia	
Spiderwort, tropical	
Thistle, Russian	
Wallflower, bushy	

Weeds Controlled	Aim EC Use Rate fl. oz. (pound active ingredient) per acre
All the weeds controlled at 1.1 fl. oz. (0.016 pound active) per acre plus the weeds listed below:	1.6 fl. oz. (0.025 pound active ingredient) per acre
Amaranth, Palmer	
Corn Spurry	
Filaree, broadleaf	
Filaree, white	
Lettuce, prickly	
Mallow, Venice (up to 2 inches tall)	
Meadowfoam	
Redmaids	
Weed List, Burndown of top growth	Aim EC Use Rate fl. oz. (pound active ingredient) per acre
Bindweed, field	1.0 – 2.0 fl. oz. (0.016 – 0.032 pound active ingredient) per acre
Burclover	
Dayflower	
Sage, lanceleaf	
Sowthistle	

***When used as directed, Aim EC will provide control of the listed weeds up to four (4) inches in height, or as specified.**