



Bayer CropScience, LP
P.O Box 12014
2 T.W Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

METHOD 240SL HERBICIDE®

For the tank mix application of Method 240SL Herbicide plus triclopyr amine or triclopyr ester for control of mesquite, huisache, and other susceptible brush species with IPT foliar leaf spray applications in non-crop sites.

EPA Reg. No.

Method 240SL: 432-1565

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW MEXICO

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

METHOD® 240SL herbicide can be tank mixed with triclopyr amine or triclopyr ester for control of mesquite, huisache, or other susceptible brush species through individual plant treatment (IPT) foliar leaf sprays in labeled non-crop sites including airports, highway, railroad, pipeline and utility rights-of-way, sewage disposal areas, industrial areas, such as electrical substations, or rail yards, farmyards, fuel storage areas, fence rows, non-irrigation ditch banks, barrier strips, lumberyards, pumping stations and tank farms, restoration areas, wildlife management areas, wildlife openings, and wildlife habitats.

Species:

Honey mesquite, western honey mesquite, huisache, whitebrush, tasajillo, lotebush (spp), bumelia, flame leaf sumac, red bud, yucca (spp), Texas persimmon, catclaw acacia, catclaw mimosa, Xanthoxylum (spp) (also known as tickle tongue, toothache tree, or prickly ash), guajillo, mountain laurel, blackbrush, coyotillo, agarito, and other susceptible species.

Mixing:

Add 64 fl oz of METHOD 240 SL herbicide and 85 fl oz/A triclopyr amine (3 lb ai/gallon formulation) OR 64 fl oz of triclopyr ester (4 lb ai/gallon formulation) and 64 fl oz of MSO adjuvant in enough water to form 100 gallons of total spray. This mixture should be sprayed over an area of at least 4 acres as not to exceed the maximum application rate of METHOD 240 SL herbicide of 18 fl oz/A.

Application Timing:

For mesquite, apply in the late-spring to mid-summer. For huisache, apply from late-summer into fall. Make applications when foliage is healthy, mature with a dark green color and when soil temperature is at least 75 degrees F at a depth of 12 inches. Control results are best when applications are made with moderate to high soil moisture and with uniform healthy foliage.

Application:

Apply herbicide mixture with a backpack, knapsack, or ATV sprayer using low pressure and an adjustable hollow cone nozzle (i.e. Conejet 5500-X6 or X8). For best results, adjust spray pressure and nozzle to deliver a coarse spray with large droplets. Thoroughly spray all the foliage of the targeted plants but not to the point of runoff.



Bayer CropScience, LP
P.O Box 12014
2 T.W Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

METHOD 240SL HERBICIDE®

For the tank mix application of Method 240SL Herbicide plus triclopyr amine or triclopyr ester for control of mesquite, huisache, and other susceptible brush species with IPT foliar leaf spray applications in non-crop sites.

EPA Reg. No.
Method 240SL: 432-1565

FIFRA 2(ee) Recommendation

IMPORTANT PRECAUTIONS:

Vegetation in close proximity to treated plants, contacted by herbicide spray, or whose root zone is contacted with herbicide spray may show injury and/or die following treatment. Natural vegetation restoration in this zone will occur over the next one to two growing seasons depending on environmental conditions. Do not use this product if this injury is not acceptable.

Certain species may, in particular, be sensitive to low levels of METHOD® 240SL including but not limited to, conifers (such as Douglas fir, Norway spruce, ponderosa pine and white pine), deciduous trees (such as aspen, Chinese tallow, cottonwood, honey locust, magnolia, poplar species, redbud, silver maple, and willow species) and ornamental shrubs (such as arborvitae, burning bush, crape myrtle, forsythia, hydrangea, ice plant, magnolia, purple plum and yew).

Injury or loss of desirable trees or vegetation may result if METHOD® 240SL herbicide is applied on or near desirable trees or vegetation, on areas where their roots extend, in locations where the treated soil may be washed or moved into contact with their roots, or where roots of treated plants and roots of nearby desirable trees extend into the same root zone. Consider site-specific characteristics and conditions that could contribute to unintended root zone exposure to desirable trees or vegetation. Root zone areas of desirable trees or vegetation are affected by local conditions and can extend beyond the tree canopy. If further information is needed regarding root zone area, consult appropriate state extension service, professional consultant or other qualified authority.

Injury to or loss of desirable trees or vegetation may result if equipment is drained or flushed on or near these trees or vegetation, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

IMPORTANT RESTRICTIONS:

DO NOT USE PLANT MATERIAL TREATED WITH METHOD® 240SL FOR MULCH OR COMPOST.

Do not apply this product in areas where the roots of desirable trees and/or shrubs may extend unless injury or loss can be tolerated. Root zone areas of desirable trees or vegetation are affected by local conditions and can extend well beyond the tree canopy.

Do not apply this product if site-specific characteristics and conditions exist that could contribute to movement and unintended root zone exposure to desirable trees or vegetation unless injury or loss can be tolerated.

During period of intense rainfall, applications made to roadsides or other non-crop areas, to soils saturated with water, or soils through which rainfall will not readily penetrate may result in runoff and movement of METHOD® 240SL. Do not apply METHOD® 240SL when these conditions exist.

Do not apply more than 18 fluid ounces METHOD® 240SL herbicide per acre per year on a broadcast acre basis.

. For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937)*