



GLORY 4L™  
EPA REG. NO. 66222-256

**Postemergence Directed Spray Application for Certain Weeds in Soybeans**  
**FIFRA Section 2(ee) recommendation for use in IA IL and MN**

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for GLORY 4L and on the last page of this 2(ee) before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by U.S. EPA.

**APPLICATION INFORMATION**

GLORY 4L can applied in postemergence directed sprays to weeds in soybeans fields for control of certain weeds which escape preplant or preemergence herbicide applications and for control of additional flushes of weeds that may occur after soybeans have emerged.

Postemergence directed sprays of GLORY 4L can be applied to soybeans in addition to a preemergence or preplant application of GLORY 4L according to label directions.

**Weeds Controlled:** GLORY 4L applied postemergence in soybeans to weeds as a directed spray according to directions on this label, will control the following at rates shown (broadcast basis) when grasses and common ragweed are less than 1 inch tall and other broadleaves are less than 3 inches tall:

Rate of Glory 4L	
6.3 - 9.5 oz/Acre	6.3 – 12.6 oz/Acre
Carpetweed ( <i>Mollugo verticillata</i> ) Cocklebur ( <i>Xanthium pensylvanicum</i> ) Crabgrass ( <i>Digitaria</i> spp.) Mexicanweed ( <i>Caperonia castaniifolia</i> ) Pigweeds ( <i>Amaranthus</i> spp.) Purslane ( <i>Portulaca oleracea</i> ) Velvetleaf ( <i>Abutilion theophrasti</i> ) Waterhemp	<b>Partial control of:</b> Horsenettle ( <i>Solanum</i> spp.) Morningglory species ( <i>Ipomoea</i> spp.) Ragweed, common ( <i>Ambrosia artemisiifolia</i> ) Spotted spurge ( <i>Euphorbia maculata</i> )

## **RECOMMENDED APPLICATIONS**

Apply proper dosage using 10 to 40 gallons of water per acre as a directed spray in a 6- to 8-inch band on each side of the row after soybeans are 8 inches tall and before broadleaf weeds are 3 inches tall and before grasses and common ragweed are 1 inch tall. For best results, the spray must cover weed foliage with minimum or no contact with soybean foliage. Add a nonionic surfactant to the spray mixture to obtain better wetting of weed leaf surfaces. To determine the correct dosage of GLORY 4L for a band application, see "Band Application" directions in the Section 3 GLORY 4L product label in the "Product Information" section.

## **RESTRICTIONS (Directed Postemergence):**

- Do not apply directly to soybeans or serious crop injury will occur.
- Do not allow spray to contact more than the lower 1/4 to 1/3 of soybean plants. Soybean leaves contacted by the spray will be killed.
- Do not apply GLORY 4L postemergence to sensitive soybean varieties. See SPECIAL PRECAUTIONS in main GLORY 4L label and refer to soybean seed provider and university information regarding the most current list of sensitive varieties.
- Do not apply under weather conditions which favor drift. To avoid injury to other crops or desirable plants from spray drift, sprayer pressure must not exceed 30 psi and the sprayer must be fitted with nozzles no smaller than 8002-T-Jet (or equivalent).
- Do not feed or graze green soybean vines.
- Do not harvest soybeans or use dry soybean vines for feed or forage within 70 days of last application.

## **LIMITATION OF WARRANTY AND LIABILITY**

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **Conditions, Disclaimer of Warranties and Limitations of Liability**.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc.d/b/a ADAMA. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ADAMA makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of ADAMA is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ADAMA disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at ADAMA's election, the replacement of product.

Glory is a registered trademark of the ADAMA Group Company.

020515 – 1.0