

CHASER ULTRA 2

SELECTIVE HERBICIDE

A PREMIUM THREE-WAY POST-EMERGENT SELECTIVE BROADLEAF HERBICIDE CONTAINING MCPA, FLUROXYPYR AND DICHLORPROP-p. FOR USE ON GOLF COURSES, PARKS, ORNAMENTAL TURF LAWNS, ALSO FOR USE ON SOD FARMS.

CONTROLS DANDELIONS, CHICKWEEDS, PLANTAINS, OXALIS, SPURGE AND MANY OTHER BROADLEAF WEEDS, SOME OF WHICH ARE LISTED ON THIS LABEL.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	. 41.60%
1-Methylheptyl Ester of Fluroxypyr: ((4-amino-3-5-dichloro-	
6-fluoro-2-pyridinyl)oxy) acetic Acid, 1-methylheptyl Ester**	4.90%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***	
OTHER INGREDIENTS:	. 45.40%
TOTAL	100.00%

Isomer Specific Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid 33.97%, 3.20 lbs./gal.

**((4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic Acid 3.40%, 0.32 lbs./gal.

***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid 6.79%, 0.64 lbs./gal.

CONTAINS THE SINGLE ISOMER FORM OF DICHLORPROP-p.

INTENDED FOR USE BY PROFESSIONAL TURF MAINTENANCE
PERSONNEL, LANDSCAPING OR COMMERCIAL APPLICATORS ONLY.
NOT INTENDED FOR SALE TO OR USE BY HOMEOWNERS.
DO NOT SELL, DISTRIBUTE OR USE THE PRODUCT IN NASSAU AND SUFFOLK COUNTY IN NEW YORK.

KEEP OUT OF REACH OF CHILDREN DANGER—PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID				
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. 			
	 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. 			
	 Call a poison control center or doctor for treatment advice. 			
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. 			
	 Have person sip a glass of water if able to swallow. 			
	 Do not induce vomiting unless told to do so by the poison control center or doctor. 			
	 Do not give anything by mouth to an unconscious person. 			
If on skin	• Take off contaminated clothing.			
or clothing:	 Rinse skin immediately with plenty of water for 15 to 20 minutes. 			
	Call a poison control center or doctor for treatment advice.			
HOT LINE NUMBER				

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.**

Note to Physician: Probable mucosal damage may contraindicate use of gastric lavage.

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA REG. NO. 34704-927 EPA EST. NO. 228-IL-1 NET CONTENTS 1 GAL. (3.78 L)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER—PELIGRO

Corrosive, causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear protective eyewear such as goggles, face shield, or safety glasses.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, and other handlers must wear the following:

- Goggles or face shield,
- · Long-sleeved shirt and long pants,
- · Chemical-resistant gloves, and
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

ENGINEERING CONTROLS STATEMENTS: If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide may adversely affect non-target plants: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

MCPA has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This product should be used strictly in accordance with the drift and runoff precautions on this label in order to minimize off-site exposure.

Under some conditions, this product may have a potential to runoff to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no-till, limited till and contour plowing; these methods also reduce pesticide runoff. Where feasible, use application techniques such as T-banding, and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetation filter strips is recommended along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where runoff could occur to minimize water runoff.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and for high temperatures.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is as follows:

- Coveralls,
- Shoes plus socks,
- Chemical-resistant gloves made of any waterproof material, and
- Protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees* and Roughs). Also for use on Sod Farms.
*Excluding Bentgrass Tees.

USE PRECAUTIONS

Aerial application of this product is prohibited.

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result. Do not use on Carpetgrass, Centipede, St. Augustine, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90°F. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Creeping jenny Annual yellow sweet clover Crimson clover Artichoke Croton Aster Cudweed Austrian fieldcress Curly dock Curly indigo Bedstraw Beggartick Dandelion Biden Dead nettle Bindweed Dock Bird vetch Dogbane Bitterweed Dogfennel Elderberry Bitter wintercress Black-eved Susan **English daisy** Black medic Fall dandelion Black mustard False flax Black-seed plantain False sunflower Blessed thistle Fiddleneck Blue lettuce Field bindweed Blue vervain Field pansy Box elder Flea bane (daisy) Bracted plantain Flixweed Florida betony **Brassbuttons** Florida pusley Bristly oxtongue Broadleaf dock Frenchweed Broadleaf plantain Galinsoga Broomweed Garlic mustard Buckhorn Goathead Buckhorn plantain Goatsbeard **Bulbous** buttercup Goldenrod Bull nettle Ground ivv Gumweed Bull thistle Hairy bittercress Burdock Hairy fleabane Burning nettle Bur ragweed Hawkweed Burweed Healall **Buttercup** Heartleaf drymary Canada thistle Hedge bindweed Carolina geranium Hedge mustard Carpetweed Hemp Catchweed bedstraw Henbit Catnip Hoary cress Hoary plantain Catsear Chickweed Hoary vervain Chicory Honeysuckle Cinquefoil Hop clover Clover Horsenettle Cockle Horsetail Cocklebur Indiana mallow Coffeebean Ironweed Coffeeweed Jewelweed Common chickweed Jimsonweed

Common mullein

Corn Chamomile

Common sowthistle

Lambsquarters Roundleafed marigold Lespedeza Rush Locoweed Russian pigweed Russian thistle Lupine Mallow St. Johnswort Marshelder Scarlet pimpernel Matchweed Scotch thistle Sheep sorrel Mexicanweed Milk vetch Shepherdspurse Milkweed bloodflower Slender plantain Morningglory Smallflower galinsoga Mugwort Smartweed Mousear chickweed Smooth dock Musk thistle Smooth pigweed Mustard Sneezeweed Narrowleaf plantain Southern wild rose Narrowleaf vetch Sowthistle Nettle Spanishneedle Orange hawkweed **O**xalis **Spatterdock** Oxeve daisy Speedwell Parsley-piert Spiny amaranth Parsnip Spiny cocklebur Pearlwort Spotted catsear Pennycress Spotted knapweed Pennywort Spotted spurge Peppergrass Spurge Pepperweed Spurweed Piaweed Stinging nettle Pineywoods bedstraw Stinkweed Plains coreopsis Stitchwort Plantain Strawberry clover Poison hemlock Poison ivv Sumac Poison oak Sunflower Pokeweed Sweet clover Poorioe Tall nettle Povertyweed Tall vervain Prickly lettuce Tansy mustard Prickly sida Tansy ragwort Primrose **Tanweed** Prostrate knotweed **Tarweed** Prostrate pigweed Thistle Prostrate spurge Tick trefoil Prostrate vervain **Toadflax Puncturevine** Purslane Trailing crownvetch Ragweed Tumble mustard Red clover Tumble pigweed Redroot pigweed **Tumbleweed** Red sorrel

Velvetleaf

Veronica

Venice mallow

Redstem filaree

Rough fleabane

Rough cinquefoil

Kochia

Knawel

Knotweed

Vervain	Wild aster	Wild parsnip	Woolly morningglory
Vetch	Wild buckwheat	Wild radish	Woolly plantain
Virginia buttonweed	Wild carrot	Wild rape	Wormseed
Virginia creeper	Wild four-o'-clock	Wild strawberry	Yarrow
Virginia pepperweed	Wild garlic	Wild sweet potato	Yellow rocket
Wavyleaf bullthistle	Wild geranium	Wild vetch	Yellowflower
Western clematis	Wild lettuce	Willow	pepperweed
Western salsify	Wild marigold	Witchweed	Other broadleaf weeds.

White clover Wild mustard Woodsorrel White mustard Wild onion Woolly croton

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. groundboom, sprayer) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size: Use only medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles.

Wind Speed: Do not apply at wind speeds greater than 10 mph.

Temperature Inversions: If applying at wind speeds less than 3 mph, the applicator must determine if (1) conditions of temperature inversion exist, or (2) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Additional Requirements for Groundboom Application: Do not apply with a nozzle height greater than four feet above the target site.

MIXING INSTRUCTIONS

Fill spray tank with water. Start agitation and slowly add the listed amount of concentrate. Maintain continuous agitation after mixing and during application. If this product is allowed to stand in tank for extended periods of time, some separation may occur. Reagitate before use.

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Ryegrass, Bentgrass (excluding golf course greens and tees), Bahia, Bermudagrass and Zoysia grass.

For broadcast treatments:

Apply a maximum of 0.75 lb. ae 2,4-DP-p per acre per application. Do not apply more than 2 applications per year with a minimum of 30 days between applications.

For spot treatments:

Apply a maximum of 2.0 lbs. ae 2,4-DP-p per acre per application. A minimum of 30 days between applications is required. Broadcast application is prohibited at this use rate.

Spot treatment is defined as a treatment area no greater than 1,000 square feet per acre.

Lower Volume Equipment: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

NOTE: For all grasses (1) avoid double coverage resulting from overlap unless temporary turf injury can be tolerated and (2) use reduced rates if grass is stressed from heat, drought, etc.

COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

- 1. Pour 18 ounces of water into a quart iar.
- 2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
- 3. Add 1 ounce of this product.
- 4. Close jar and shake well.
- 5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
- 6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
- 7. Compatibility may be improved by the use of a compatibility agent. Loveland Products, Inc. suggests using E-Z Mix. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
- 8. If the mixture does not separate, gel, foam, or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 25°F. Protect product from freezing. If allowed to freeze, remix well before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons: If rinsate cannot be used, follow pesticide disposal instructions. If not triple rinsed or pressure rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state and federal regulations. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 5 gallons and less than 56 gallons: If rinsate cannot be used, follow pesticide disposal instructions. If not triple rinsed or pressure rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state and federal regulations. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 56 gallons: If rinsate cannot be used, follow pesticide disposal instructions. If not triple rinsed or pressure rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local,

Storage & Disposal cont'd.:

state and federal regulations. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Chaser is a registered trademark of Loveland Products, Inc.

FORMULATED FOR LOVELAND PRODUCTS, INC. P.O. BOX 1286, GREELEY, COLORADO 80632-1286