



Directions for Use on Almonds to Control Scab, Alternaria Leaf Spot, and Suppress Hull Rot

For use in California

PRODUCT BULLETIN

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

APPLICATION DIRECTIONS

Crop	Diseases	Use Rate Fl. Oz.	Instructions
		Product Per Acre	
Almond	Scab	12.5 to 15 fl oz	Application Instructions:
	(Cladosporium		Initiate applications when conditions are favorable for disease
	carpophilim)		development and continue on a 7-14 day interval. If disease pressure
	Alternaria leaf spot		is severe use the higher rate and shortest interval.
	(Alternaria		
	alternate)		Apply this product in sufficient water to obtain adequate coverage of
	,		the foliage. Spray volume will usually be 50 to 100 gallons per acre for dilute sprays and 5 to 10 gallons per acre for aerial sprays. For
	Suppression:		aerial applications, apply this product in a minimum of 5 gallons of
	Hull rot		water per acre. For dilute sprays, if higher spray volumes are desired
	(Monilinia spp., Rhizopus spp.)		for improved coverage, do not exceed the maximum rate of 15 fl oz of
	TitiiZopus spp.)	Типгориз эрр.)	product per acre.
			Resistance Management:
			Do not make more than 2 sequential applications of this product or
			other Group 3 or Group 7 containing fungicide before rotating to a
			fungicide with a different mode of action. Do not apply a third
			application of this product within 14 days of the second application.
			Restrictions:
			Do Not apply more than 63 fl oz/A/year.
			The pre-harvest interval (PHI) for this crop is 40 days.
			Do Not apply more than the maximum yearly rate from any
			combination of products containing Isofetamid (1.77 lb. a.i. per acre
			per year) or Tebuconazole (0.9 lb. a.i. per acre per year) to this crop.

For **24-Hour Medical Emergency Assistance** (Human or Animal) Call 1-888-484-7546. For **Chemical Emergency**, Spill, Leak, Fire or Accident, Call CHEMTREC 1-800-424-9300.

Read and follow all applicable directions for use, precautions and limitations on the product label attached to the $FERVENT^{TM}$ 475SC Fungicide container.



Summit Agro USA, LLC 240 Leigh Farm Road, Suite 215 Durham, NC 27707