

FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Starane® Ultra

EPA Reg. No. 62719-577

24(c) Special Local Need Registration EPA SLN No. WA-210006

(FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON)

Expiration Date: This label for Starane Ultra expires and must not be distributed or used in accordance with this SLN registration after December 31, 2025

Control of Kochia, Volunteer Potatoes, and Other Susceptible Broadleaf Weeds in Dry Bulb Onions

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Starane® Ultra herbicide before applying. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- Use of Starane Ultra according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Starane Ultra.

Directions for Use

Starane® Ultra herbicide may be applied for postemergence control of kochia, volunteer potatoes, and other susceptible broadleaf weeds in dry bulb onions using ground or aerial application equipment. Refer to the product label for Starane Ultra for a complete list of weeds controlled. Follow all mixing and application instructions in the Product Information section of the product label for Starane Ultra.

Rate and Application Timing: Apply 0.35 to 0.7 pint per acre of Starane Ultra as a broadcast postemergence treatment. Volunteer potatoes and other susceptible target weeds should be from 4 to 8 inches tall for optimum control. Kochia should be from 1-4 inches tall for optimum control.

Broadcast (over-the-top) application may be made to dry bulb onions from the 2 true leaf stage through the 6-leaf stage. Do not apply as a broadcast over-the-top spray after the 6-leaf stage of growth. Application to dry bulb onions beyond the 6-leaf stage must be made as a directed spray using drop nozzles (see crop injury warning below). Tank mix combinations with other herbicides registered for use in dry bulb onions may result in unacceptable crop injury. Adjuvants are not recommended with Starane Ultra applications in dry bulb onions.

Sequential Applications: To control heavy populations or successive flushes of volunteer potatoes, or other susceptible broadleaf weeds, two postemergence applications can be made on a 10- to 14-day re-treatment interval. Do not make more than two applications per season.

Crop Injury Warning: Crop injury (such as but not limited to leaf twisting) may occur with some dry bulb onion cultivars when Starane Ultra is applied as a broadcast treatment, especially when applications are made to larger dry bulb onions. Do not use Starane Ultra if the risk of injury is unacceptable.

Specific Use Restrictions:

- Do not harvest treated forage or allow livestock to graze treated areas within 7 days of application.
- **Preharvest Interval:** Do not apply within 42 days of dry bulb onion harvest.
- Do not make more than 2 applications per season.
- Do not apply more than a total of 0.7 pint per acre of Starane Ultra per crop season.
- **Plant-Back Restriction:** Plant only labeled crops within 120 days of application.
- **Chemigation:** Do not apply through any type of irrigation system.
- Do not apply this product when furrow irrigation is running. Treated fields should be managed to avoid water runoff for at least 6 hours after application.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site (<https://agr.wa.gov/departments/pesticides-and-fertilizers/pesticides/waste-pesticide>). Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

Conditions and Risks of Use for Special Local Need

USE OF Starane® Ultra HERBICIDE (THE “PRODUCT”) ON ONIONS (THE “CROP”) FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP.

This Product is available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Supplemental Labeling, you acknowledge and accept that to the extent permitted by law:

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Corteva Agriscience does not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental labeling;
- (3) Corteva Agriscience does not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) these Conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Corteva Agriscience, or its respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

These Conditions and Risks of Use for Special Local Need are required by Corteva Agriscience and not specified by the US EPA or the State of Washington.

TM[®] Trademarks of Corteva Agriscience and its affiliated companies

Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268

R357-041
Accepted: 05/24/21
Initial Printing