

# ZinManMax™

**A zinc, manganese, molybdenum and boron true solution with PAC\* chelation technology**

## GUARANTEED ANALYSIS:

Boron (B) . . . . .	0.5%
Manganese (Mn) . . . . .	2.7%
2.7% Chelated Manganese	
Molybdenum (Mo) . . . . .	0.25%
Zinc (Zn) . . . . .	5.1%
5.1% Chelated Zinc	

Derived from boric acid, manganese sulfate, sodium molybdate and zinc oxide chelated with PolyAldoCarbosate®

**For use as a fertilizer in foliar, fertigation and soil applications to supplement micronutrient levels on agricultural crops.**

\*PAC refers to NutriAg's "PolyAldoCarbosate" Technology

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

## DIRECTIONS FOR USE (Use only as Directed):

- Exercise caution with foliar applications in extreme weather conditions such as high temperatures and poor drying conditions.
- Aerial application – minimum of 5 gallons of water/acre; Dilute sprays – minimum of 20 gallons of water/acre. Apply with sufficient water to deliver spray uniformly on leaf surfaces. Higher volumes of water should be used for higher rates of fertilizer.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- ZinManMax is compatible with most agricultural chemicals and nutrients. Not compatible with aluminum equipment. Should be applied in conjunction with tissue and/or soil analysis.
- Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions.
- Consult a NutriAg representative for specific crop recommendations, application timings and necessity of its use.**

**KEEP OUT OF REACH OF CHILDREN.**

**Net Contents:**

**Net Weight:**

**Net Weight/  
Gallon:**

**Batch No.:**

## SUGGESTED RATES

**FOLIAR RATES:** For minor - moderate deficiency apply 1 qt/ac. For severe deficiency & post-harvest trees and vines apply 1.5-4 qt/ac. Multiple applications may be required.

**FERTIGATION RATES:** The number and frequency of applications determine application rates. Apply 1-3 qt/ac/application. Apply no more than 9 qt/season. Always apply toward the end of an irrigation cycle.

**SOIL RATES:** This product is compatible with many liquid fertilizers. Fertilizers vary in analysis and quality, and should be jar tested with this product before use. Apply up to 3 quarts/acre at planting or as a side dress.

**CROPS: Forage Crops:** Alfalfa, Clover, Hay Crops; **Field Crops:** Cereals, Corn, Cotton, Sorghum, Sugar cane; **Legumes:** Dry bean, Lentil, Peanut, Pea, Soybean; **Root Vegetables (including, but not limited to):** Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip; **Leafy Vegetables (including but not limited to):** Bok Choy, Celery, Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach; **Other Vegetables:** Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato; **Berry Crops:** Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry; **Tree Fruit (including but not limited to):** Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince; **Other Fruit:** Citrus (all), Fig, Kiwi, Melons; **Nut Trees (including but not limited to):** Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut; **Grapes:** Currant, Raisin, Table, Wine; **Other Crops:** Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower

**ATTENTION: This fertilizing material contains boron. The use of this fertilizing material on any crops other than those recommended may result in serious injury to crop(s).**

**ATTENTION: The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.**

**Warranty Statement:** Sellers guarantee shall be limited to the terms set out on the label. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg. NutriAg is not liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.



## DANGER

Causes serious eye damage. Causes skin irritation. Suspected of causing genetic defects. May cause damage to organs through prolonged or repeated exposure. Contains 18-22% of ingredients of unknown toxicity (Inhalation and Dermal). Contains 2-3% of ingredients of unknown toxicity (Oral).

Do not breathe mist, vapours, spray. Wash hands and skin thoroughly after handling. Wear protective gloves. Wear eye protection/face protection.

**IF ON SKIN:** Wash with plenty of water. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER/doctor. Get medical advice/attention if you feel unwell. Specific treatment (see supplemental first aid instruction on this label). If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse.

Dispose of contents/container in accordance with local, regional, national and international regulations.

Rev: 121917US-CA/US

**NutriAg Ltd.**

39 Gail Grove, Toronto, ON, Canada, M9M 1M5. Tel: (416) 636-1555

**[nutriag.com](http://nutriag.com)**

**NutriAg**

INNOVATORS IN THE FIELD