

# BIOMIN<sup>®</sup> BORON<sup>POWDER</sup>



## Chelated Minerals for Soil and Foliar Application

### PRODUCT INFORMATION

**BIOMIN BORON** - Contains chelated mineral for soil and/or foliar applications. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct BORON deficiency and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizers.

**BIOMIN BORON** may be applied to FRUIT TREES, FIELD CROPS, GREENHOUSE CROPS, VEGETABLES, POTATOES, GRAPES, CITRUS, TURF and ORNAMENTAL GARDENS, etc...

### DIRECTIONS AND GENERAL RECOMMENDATIONS:

#### SOIL APPLICATION

**BIOMIN BORON** may be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1.5-5.0 pounds (1.0-5.5 Kg/Ha) per acre depending on crop, soil conditions and application methods. For maintenance, 1.5 pounds (1.5 Kg/Ha) per acre are usually recommended.

#### FOLIAR APPLICATION

**FIELD CROPS:** Apply 1.0-1.5 pounds (1.0-1.5 Kg/Ha) of **BIOMIN BORON** per acre when crops are 6-12 inches (15-30 cm) high or when deficiency symptoms appear. Repeat application 2-4 weeks later.

**VEGETABLES and POTATOES:** Apply 1.0-1.5 pounds (1.0-1.5 Kg/Ha) of **BIOMIN BORON** per acre at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later.

### GUARANTEED ANALYSIS

	TOTAL PRODUCT BASIS %W/W
Boron (B)	15.0
Derived from sodium borate and glycine	

### PREVENTS AND CORRECTS BORON DEFICIENCY

**CAUTION** - This product contains boron. Do not apply to boron sensitive crops. Do not apply to crops that will be grazed by ruminant animals. Contact your local county agency or field consultant for further information.

### KEEP OUT OF REACH OF CHILDREN

Manufactured by:  
JH BIOTECH, INC.  
P.O. Box 3538  
Ventura, CA 93006  
U.S.A.

Lot #:  
U.S. Pat.# 5,504,055

**GREENHOUSE CROPS:** Apply 1.0-1.5 pounds (1.0-1.5 Kg/Ha) of **BIOMIN BORON** per acre 2-3 weeks after emergence or transplanting, or when magnesium deficiencies first appear. Repeat the application every one or two weeks.

**FRUIT TREES AND GRAPES:** Apply 1.0-1.5 pounds (1.0-1.5 Kg/Ha) of **BIOMIN BORON** per acre in early spring when new leaves emerge or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

**BIOMIN BORON** is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 100 times of water. For best results and more specific recommendations, contact your nearest distributor.

### LIMITED WARRANTY

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

### NET CONTENTS

- |  |  |
|--|--|
| <input type="checkbox"/> 2.5 pounds (1 Kg) | <input type="checkbox"/> 5 pounds (2.5 Kg) |
| <input type="checkbox"/> 25 pounds (10 Kg) | <input type="checkbox"/> 55 pounds (25 Kg) |

MANUFACTURING DATE:

0407-D-A-1

# JH BIOTECH, INC.