### SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1070A-S4A 0918

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

FLUAZIFOP-P-BUTYL GROUP 1 HERBICIDE

#### Fusilade® DX Herbicide

This supplemental label expires on 08/31/2020 and must not be used or distributed after this date.

Active Ingredient:

Fluazifop-P-butyl

Butyl ( <i>R</i> )-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate*	24.5%
Other Ingredients:	75.5%
Total:	100.0%

Fusilade® DX Herbicide formulation is an emulsifiable concentrate (EC).

#### KEEP OUT OF REACH OF CHILDREN.

# **CAUTION**

EPA Reg. No. 100-1070

All applicable directions, restrictions and precautions on the Fusilade DX Herbicide (EPA Reg. No. 100-1070) label are to be followed. Before using Fusilade DX Herbicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Use of Fusilade DX Herbicide according to this labeling is subject to the limitations imposed by the label affixed to the container for Fusilade DX Herbicide (EPA Reg. No. 100-1070).



<sup>\*</sup>Fusilade DX Herbicide contains 2 pounds (+) isomer (fluazifop-P-butyl) per gallon.

# **DIRECTIONS FOR USE**

### **Fusilade DX Rate Conversion Table**

FI oz product/acre	Lb ai fluazifop/acre
4	0.062
6	0.094
8	0.125
10	0.156
12	0.188
16	0.250
24	0.375
30	0.469
32	0.500
48	0.750
72	1.125

# FINE FESCUE GRASSES GROWN FOR SEED (IDAHO, OREGON AND WASHINGTON ONLY)

- Do not apply a total of more than 48 fl oz of Fusilade DX Herbicide per acre per vear.
- Do not apply more than 16 fl oz of Fusilade DX Herbicide per acre per application.
- Do not apply more than 3 applications per year.
- Maintain a minimum of 14 days between applications.
- Do not harvest forage within 14 days of application.

# SPECIAL USE DIRECTIONS FOR FINE FESCUE GRASSES GROWN FOR SEED IN IDAHO, OREGON AND WASHINGTON

For control of downy brome, quackgrass, bentgrass and volunteer cereals in fine fescue grasses (such as Chewings, Creeping Red and Hard Fescue varieties) grown for seed. **NOT FOR USE ON TALL FESCUE**. Refer to Tables 2 and 4 for a complete listing of grass weeds controlled. Refer to the Spray Additives section for spray additives.

## **Application Rate and Timing**

**Fall Application** - Apply Fusilade DX Herbicide at 8 fl oz per acre to actively growing downy brome in the 2 to 4 inch stage. Apply Fusilade DX Herbicide at 12 to 16 fl oz per acre to actively growing quackgrass or bentgrass in the 6 to 10 inch stage and make a second application of 12 to 16 fl oz per acre before regrowth exceeds 10 inches.

**Spring Application** - Apply Fusilade DX Herbicide at 8 to 16 fl oz per acre to actively growing grasses. Downy brome in the 2 to 4 inch stage can be controlled by the lower rate but larger, established grasses (4 to 6 inches) may need the higher rate for adequate control. Apply Fusilade DX Herbicide at 12 to 16 fl oz per acre when quackgrass and bentgrass is at the 6 to 10 inch stage and make a second application of 12 to 16 fl oz per acre before regrowth exceeds 10 inches.

### Restrictions:

- Refer to the CROP SPECIFIC RESTRICTIONS section for additional information.
- Do not apply Fusilade DX Herbicide after fescue grass seedhead develops into the boot stage.

### **Precautions:**

• For use on fine fescues (Chewings, Creeping Red and Hard Fescue varieties) grown for seed. Other types of fescues may be injured by Fusilade DX Herbicide.

Fusilade® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2018 Syngenta