



**Dow AgroSciences**

**Dow AgroSciences LLC**

**9330 Zionsville Road**

**Indianapolis, IN 46268-1054 USA**

# **FIFRA Section 24(c) Special Local Need (SLN) Labeling**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON**

**Spray Interval (10-14 days) in Hops for Control of Powdery Mildew**

**Quintec®**

**EPA Reg. No. 62719-375**

**EPA SLN No. WA-180001**

**This label for Quintec expires and must not be distributed or used in accordance with this SLN registration after December 31, 2022.**

## **ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

## **Directions for Use**

Refer to product label for General Use Precautions, Mixing and Application instructions.

**Maximum Application Rate:** Do not apply more than 8.2 fl oz of Quintec per acre per application or more than 33 fl oz of Quintec per acre per calendar year.

**Minimum Spray Interval:** 10 – 14 days. Use the shorter interval within the range for severe or threatening disease conditions.

## **RESTRICTIONS**

- **Preharvest Interval:** Do not apply within 21 days of harvest.
- Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.
- For use in accordance with this SLN label do not apply this product through any type of irrigation system.

## **WA Aquatic Advisory**

This pesticide is toxic to fish and aquatic invertebrates. Quintec should not be used under this SLN label where impact of listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries (NOAA) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

**WSDA Container Disposal Guidance:** Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx>. Cleaned containers may also be disposed of in a sanitary landfill if permitted by the county. Burning is not a legal method of container disposal in Washington.

®™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R136-076

Approved: 02/14/18

Initial printing.