

# Scotts Fertilizer 0-0-7 with

# HALTS II

---

For preemergence control of grass and broadleaf weeds in established turf grasses  
(excluding golf course putting greens) and lawns.

**ACTIVE INGREDIENT:**

Prodiamine: [N <sup>3</sup> ,N <sup>3</sup> -Di-n-propyl-2, 4-dinitro-6-(trifluoromethyl)-m-phenylenediamine].....	0.28%
OTHER INGREDIENTS:.....	99.72%
TOTAL.....	100.00%

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

See label for additional precautionary statements.

**NET WEIGHT: 50 POUNDS (22.7 kg.)**

## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION :** Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

#### **FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

**IF SWALLOWED:** Call a poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-561-6584 for emergency medical treatment information.

### **ENVIRONMENTAL HAZARDS**

This pesticide may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **APPLICATION PRECAUTIONS**

Not for use on turf or plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on turf or plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

**NOTE:** Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

## PRODUCT INFORMATION

This product is a turf fertilizer and a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), and lawns, landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development.

### ESTABLISHED TURF

**Application Directions:** Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation prior to weed seed germination and within 14 days of application.

**Use Precautions:** The following precautions apply to the use of this product in turf grasses and lawns.

1. Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
2. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).
3. Do not cut (harvest) treated sod before 120 days after application. Do not apply newly set sod until the following year.
4. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury.
5. Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
6. Do not apply to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

**Rates of Application:** This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential application may be made as long as the total amount of product applied does not exceed the maximum annual application rates listed for each turf species. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

**Maximum Annual Rates:** This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not apply more than 535 lbs. of this product per calendar year. Use higher rates of product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum yearly rate listed.

<b>MAXIMUM APPLICATION RATE/CALENDAR YEAR (by Turf Grass Species)</b>		
<b>Turf Species</b>	<b>Lbs. Product/Acre</b>	<b>Lbs. a.i./Acre</b>
Creeping Bentgrass	233 lbs.	0.65 lb.
Creeping Red Fescue	267 lbs.	0.75 lb.
Buffalograss Kentucky Bluegrass Perennial Ryegrass	357 lbs.	1.00 lb.
Bahiagrass <sup>1</sup> Bermudagrass <sup>2</sup> Centipedegrass <sup>1</sup> Seashore Paspalum <sup>1</sup> St. Augustinegrass Tall Fescue (including turf-Type) Zoysiagrass	535 lbs.	1.50 lb.

<sup>1</sup>These are the maximum rates per calendar year.

<sup>2</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 180 pounds/acre. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

**When To Apply After Overseeding Turf:** Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

**When To Overseed After Application:** This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results.

<b>Product Rate Lbs./Acre</b>		<b>Months Before Overwintering</b>		
<b>Lbs. Product/Acre</b>	<b>Lbs. a.i./Acre</b>	<b>North</b>	<b>Transition</b>	<b>South</b>
180	0.50	4	4	4
233	0.65	5	4	4
267	0.75	6	5	5
285	0.80	-	6	6
357	1.00	-	7	7
407	1.14	-	-	9
465	1.30	-	-	10
535	1.50	-	-	12

**Application Directions:** Apply uniformly with suitable, calibrated application equipment. Calibrate your spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. Apply this product uniformly over the area to be treated. Avoid streaking, skips and overlaps during application. Calibration of spreader may be needed if conditions of area to be treated changes and rate or speed of application are changed.

SPREADER RECOMMENDATIONS				
Spreader Settings for 1000 sq. ft.	180 lbs. product/A	267 lbs. product/A	357 lbs. product/A	535 lbs. product/A
Scott's Evergreen 18"	9-10	10-11	10-12	12-14
Scott's Speedy Brdct.	L-M	M-N	M-O	O-P
Earthway Broadcast	20-21	21-22	21-23	23-25
Earthway Drop 18"	13-14	14-15	14-16	16-17
Lesco	L-M	M-N	M-O	O-P

## LANDSCAPE ORNAMENTALS

(including established perennials and wildflower plantings)

This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 5.35 to 12.28 pounds/1,000 square feet (233 to 535 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 12.28 pounds/1,000 square feet (535 pounds/acre) per year.

**Application Timing And Information:** This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

**Tolerant Ornamental Species:** This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

### SCIENTIFIC NAME

### COMMON NAME

Abies spp.	Fir species** (Balsam, Frasier, Noble, etc.)
Abelia grandiflora	Abelia: Sherwood
Acer palmatum	Japanese Maple
A. Platanoides	Norway Maple
Achillea spp.	Yarrow: King Edward
Actindia chinensis	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Agapanthus orientalis	
Akebiaq quintata	Five-leaf or Chocolate Vine
Allium cemuum	Lady's Leek, Nodding Onion
Anemone hybrida	Japanese Anemome

Aquilegia spp.	Aquilegia: Red and Gold
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Artemisia spp.	Wormwood; Silver Mound, Castle
Aster spp.	Aster: Bonnie Blue, Purple Dome
Aster X Frikartii	
Athrium Filiz-femina	Lady Fern; Fern Lady
Aucuba japonica	Japanese Aucuba
Begonia spp.	Fibrous Begonia: Hardy Grandis
Berberis gladwynensis	Barberry
B. Julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. Thunbergil	Japanese Barberry
B. Verruculosa	Warty Barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Campanula carpatica	Tussock Bellflower, (White Clips)
Campis X Tagliabuana	Trumpet Flower, Madame Galen
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. Stolonifera	American Dogwood
Cortaderia seloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. Microphyllus	Rockspray Cotoneaster
Cretaegus spp.	Hawthorne
Crocsmia spp.	Lucifer
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma spp.	Cooperi Pink
Delphinium spp.	Larkspur: Blue Elf
Dianthus deltoides	Dianthus: Maiden Pinks "Zing"
D. gratianopolitanus	Cheddar Pink
Dodonea viscosa	Hop Bush

Echinacea purpurea	Coneflower, Purple; Magnus
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
E. japonica	Evergreen Euonymus
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
Forsythia suspensa	Weeping Forsythia
F. viridissima	Greenstem Forsythia
Gaillardia spp.	Gaillardia, Blanket Flower: "Goblin"
Gardenia jasminoides	Gardenia, Cape-Jasmine
Gaura spp.	
Gentiana dahurica	Gentian
Geranium cinereum	Cranesbill
Gladiolus spp.	Gladiolus species**
Gypsophilia repens	Baby's Breath
Hedera helix	English Ivy
Hellanthemum spp.	Sunrose
Hemerocallis spp.	Daylily: Aztec Gold, Stella de Oro, Tender Love
Hibiscus	Rose of Sharon**
Hibiscus spp.	Mallow: Disco Belle White
Hibiscus rosa-sinensis	Chinese Hibiscus
Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
H. Sieboldiana	Hosta, "Seersucker"
Houttuynia cordata var. variegata	
Hydrangea macrophylla	Bigleaf Hydrangea
Ilex Bernyi	Holly
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
I. vomitoria	Yaupon Holly, Schillings
Inula ensifolia	
Iris ensata	Sword-Leaved Iris; Jodlesong
Iris siberica	Siberian Iris; Cabernet
Iris spp.	Iris species**
Jasminium nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonnii
J. conferta	Shore Juniper; Blue Pacific
J. davurica	Parsoni
J. horizontalis	Creeping Juniper
Juglans spp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crape Myrtle
Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora
Lantana montevidensis	Weeping Lantana
Lavender spp.	Lavender; Munstead



Leonopodium alpinum	Edelweiss
Ligustrum amurense	Amur. Privet
Ligustrum sinense	Chinese Privet; Variegata
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (Waxleaf)
Lilium spp.	Lily; Jazz
Liriope muscari	Big Blue Lily, Silver Mound, Evergreen Giant
Liriope muscari var. variegata	Liriope, Variegated
Liriope spicata	Liriope, Creeping
Lobelia Cardinalis	Cardinal Flower, Indian Pink
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Lorapetalum chinense	Burgundy
Lythrum spp.	Loosestrife; Modern Pink
Magnolia spp.	Magnolia species**
Maleophora luteola	Ice Plant
Malus spp.	Crabapple*
Miscanthus sinensis	Yaku Jima, Silberfeder
Nandina domestica	Heavenly Bamboo
Narcissus spp.	Narcissus species**
Nerium oleander	Oleander
Olea europaea	Olive*
Ophiopogon japonicus	Mondo Grass
Osmanthus heterophyllus	Osmanthus (False Holly); Gulf Tide
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboreum	Sourwood
Paeonia suffruticosa	Tree Peony
Pennisetum setaceum	Fountain Grass (Dwarf)
Perovskia atriplicifolia	
Persea americana	Avocado*
Photinia Fraseri	Fraser's Photinia (Redtip)
Physotegia Virginiana	Dragonhead, False; Vivid
Picea spp.	Spruce species** (Colorado Blue, Norway, etc.)
Pieris japonica	Lily-of-the-Valley Bush
Pinus brutia	Calabrian Pine
P. canariensis	Canary Island Pine
P. Elliottii	Slash Pine
P. halepensis	Aleppo Pine
P. nigra	Austrian Pine
P. palustris	Longleaf Pine
P. radiata	Monterey Pine
P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine
P. taeda	Loblolly Pine
P. Thunbergiana	Japanese Black Pine
P. Virginiana	Virginia Pine



Pistachio spp.	Pistachio*
Pittosporum rhombifolium	Queensland Pittosporum
P. tobira	Japanese Pittosporum
Podocarpus macrophyllus	Japanese Yew
Prunus laurocerasus	English Laurel
Prunus spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
Pseudotsuga menziesii	Douglas Fir**
Pyracantha coccinea	Firethorn (Scarlet)
P. fortuneana	Firethorn
P. koidzumii	Firethorn
Pyrus spp.	Bradford Pear spp.
Quercus Shumardii	Oak, Shumard's Red
Quercus spp.	Oak species
Raphiolepis indica	Indian Hawthorne
Raphiolepis umbellata	Yedda Hawthorne
Rhododendron	'Coral Bells,' Formosa,' Hino-crimson,' PJM'
(including Azalea)	'Delaware Valley White', ' Roseum Elegans,' 'Flame Creeper', Girard Crimson', 'Wakeilebisu 'White Gumbo', 'George L. Tabor'
Rosa banksiae	Lady Bank's Rose
Rosmarinus officinalis	Rosemary*
Rudbeckia spp.	Black-Eyed Susan; Goldsturm
Santolina virens	Santolina
Saxifraga spp.	Saxifrage; Purple Dome
Scabiosa spp.	Pincushion Flower
Sedum album	Stonecrop
Sedum caucicola	Stonecrop; Lidakense
Sedum dasyphyllum	Stonecrop
Sedum spurium	Stonecrop; Dragon's Blood
Spiraea bumalda	Spirea; Anthony Waterer
Syzygium paniculatum	Australian Brushcherry
Taxus cuspidata	Japanese Yew
T. media	Yew
Teucrium spp.	Germander
Thalictrum diptercarpum	Meadow Rue
Thuja occidentalis	American Arborvitae
Trachelospermum asiaticum	Star Jasmine
Tsuga canadensis	Canada Hemlock
Tulipa spp.	Tulip species
Veronica spp.	Veronica, Speedwell; Sunny Border
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. suspensum	Arrowwood Viburnum
V. tinus	Laurustinus
V. trilobium	Cranberry Bush

V. wrightii	Leatherleaf Viburnum
Vinca major	Greater Periwinkle, Vinca
Vinca Minor	Common Periwinkle, Vinca
Vitis spp.	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet, Yucca
Y. filametosa	Adam's Needle, Yucca

\*Do not use on food producing trees, vines or plants.

\*\*Not for use on container grown plants.

## WEEDS CONTROLLED

When used in accordance with label directions, this product will provide control of the following weeds:


Barnyardgrass	Henbit	Purslane, Common
Bluegrass, Annual (Poa annua)	Itchgrass	Pusley, Florida
Carpetweed	Johnsongrass (from seed)	Rescuegrass <sup>3</sup>
Chickweed, Common	Junglerice	Shepherd's Purse <sup>2</sup>
Chickweed, Mouseear (from seed)	Knotweed	Signalgrass, Broadleaf
Crabgrass (large, smooth)	Kochia	Speedwell, Persian
Crowfootgrass	Lambsquarter, Common	Sprangletop
Cupgrass, Woolly	Lovegrass	Spurge, Prostrate
Foxtails, Annual	Panicum (Texas, fall, browntop)	Witchgrass
Goosegrass <sup>1</sup>	Pigweed	Woodsorrel, Yellow (from seed)

<sup>1</sup>In many areas, a single application of 233 to 535 pounds/acre will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 233 to 357 pounds/acre, followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

<sup>2</sup>For this weed, apply in late summer, fall or winter prior to germination.

<sup>3</sup>Suppression only.

### GERMINATION DATES



# GUARANTEED ANALYSIS

## Scotts Fertilizer 0-0-7 with Halts II

0-0-7

F643

Soluble Potash ( $K_2O$ )..... 7%

**Derived From:** potassium chloride

Chlorine (Cl), not more than ..... 4.8%

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

**FOR FLORIDA:**

**We recommend that you follow the Green Industries BMP's at:**

[http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP\\_Book\\_final.pdf](http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP_Book_final.pdf)

**FOR FLORIDA:**

**We recommend that you follow the Golf Course BMP's at:**

<http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE AND CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. **If spilled:** Sweep up and use as per label instructions or dispose of as indicated above.

NOTICE TO BUYER – Purchase of this material does not confer any rights under patents of countries outside of the United States.

## **LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of The Scotts Company or Seller. These risks can cause: ineffectiveness of the product or other unintended consequences. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

The Scotts Company warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, The SCOTTS COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THE SCOTTS COMPANY OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF THE SCOTTS COMPANY OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHER LEGAL THEORY) RESULTING FROM THE STORAGE, USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF THE SCOTTS COMPANY OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, The Scotts Company or its distributor must have prompt notice of any claim so that an immediate investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify The Scotts Company or a Scott's Company distributor or any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

**EPA Reg. No.:** 538-316

**EPA EST. NO.**

8378-IN-1<sup>k</sup>

35512-FL-2<sup>H</sup>

82757-MA-002<sup>E</sup>

84175-TX-1<sup>B</sup>

9198-OH-1<sup>M</sup>

538-OH-1<sup>S</sup>

35497-OR-001A<sup>Y</sup>

Superscript used is first letter of run code on end of container.

Manufactured by:  
The Scotts Company  
14111 Scottslawn Rd.  
Marysville, OH 43041

© 2010. Licensed by The Scotts Company LLC.  
World rights reserved  
Made in USA