



Bayer CropScience

Bayer Environmental Science, a Division of
Bayer CropScience, LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Mirage Stressgard®

EPA Reg. No. 432-1529

For use on Leaf and Sheath Spot (*Rhizoctonia zeae*),
Take-All Root Rot, and Spring Dead Spot
Expiration 05/03/2019

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF Florida, Hawaii, Georgia, South Carolina, North Carolina, Tennessee, Arkansas, Oklahoma, Mississippi, Louisiana, New Mexico and Arizona

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling. The user must refer to the federally approved label for Mirage Stressgard® and read and follow all directions for use, restrictions, and precautions. For Use on turfgrass and golf courses only.

Restrictions:

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational parks, athletic fields, athletic fields located on or next to schools, campgrounds, churches, and theme parks. Do not use on turf grown for sale or commercial use as sod. Not for residential use.

Disease Controlled	Rate of Mirage Stressgard®/1,000 ft sq & Recommended Applications
Leaf and Sheath Blight, Mini Ring (<i>Rhizoctonia zeae</i>)	1.0-2.0 fl. oz. Apply preventively or curatively when conditions are favorable for disease development. Space applications 21-28 days apart. A tank mix of Mirage Stressgard® at 1 fl. oz. per 1,000 sq. ft. and Prostar® 70WG (432-1223) at 2.2 oz. per 1,000 sq. ft. can be used when conditions of high disease pressure exist.
Take-All Root Rot, Bermudagrass Decline, Warm Season Turfgrass Decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	1.0-2.0 fl. oz. Apply preventive fungicides in the spring and fall. Make 1-2 applications prior to conditions favorable for disease development. Space applications 28 days apart. Apply fungicide in adequate water volume and water-in applications to root zone.
Spring Dead Spot (<i>Ophiosphaerella korrae</i> , <i>O. herpotricha</i> , <i>O. namar</i>)	1.0-2.0 fl. oz. Begin fungicide applications preventively in the fall when soil temperatures drop below 75° F at a 2-inch soil depth. Make 2 applications. Space applications 28 days apart. Apply fungicide in adequate water volume and water-in applications to root zone.

When to apply: Apply fungicides preventively unless otherwise indicated. Make applications prior to conditions favorable for disease development. Interval between apps: 28 days unless otherwise noted.

Application Directions:

Maximum use is limited to six applications or 6.5 fl. oz. of Mirage Stressgard® per 1000 sq. ft. per year. For applications on ultra-dwarf bermudagrass only use the low label rate of 1 fl. oz. per 1000 sq. ft.

**For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information, Call 1-866-99BAYER (1-866-992-2937)**

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*