

# Supplemental Labeling



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Starane<sup>®</sup> Ultra

EPA Reg. No. 62719-577

### Control of Kochia, Volunteer Potatoes, and Other Susceptible Broadleaf Weeds in Dry Bulb Onions

This supplemental label expires April 1, 2020, and must not be used or distributed after this date.

(For Distribution and Use Only in the States of Colorado, Idaho, Oregon, and Washington)

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Starane<sup>®</sup> Ultra herbicide before applying. Follow all applicable directions, restrictions, PPE requirements and precautions on the label affixed to the container for Starane Ultra.
- Use of Starane Ultra according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Starane Ultra.

## WARNING

## AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Directions for Use

Starane<sup>®</sup> Ultra may be applied for postemergence control of kochia, volunteer potatoes, and other susceptible broadleaf weeds in dry bulb onions using ground or aerial application equipment. Refer to the product label for Starane Ultra for a complete list of weeds controlled. Follow all mixing and application instructions in the Product Information section of the product label for Starane Ultra.

**Rate and Application Timing:** Apply 0.35 pint Starane Ultra as a broadcast postemergence treatment. Volunteer potatoes, kochia, and other susceptible target weeds should be from 4 to 8 inches tall for optimum control.

Broadcast (over-the-top) application may be made to dry bulb onions from the 2 true leaf stage through the 6-leaf stage. Application to dry bulb onions beyond the 6-leaf stage should be made as a directed spray using drop nozzles (see crop injury warning below). Do not apply as a broadcast over-the-top spray after the 6-leaf stage of growth. Tank mix combinations with other herbicides registered for use in dry bulb onions may result in unacceptable crop injury. Adjuvants are not recommended with Starane Ultra applications in dry bulb onions.

**Sequential Applications:** To control heavy populations or successive flushes of kochia, volunteer potatoes, or other susceptible broadleaf weeds, two postemergence applications can be made on a 10- to 14-day re-treatment interval. Do not make more than two applications per season.

**Crop Injury Warning:** Crop injury such as but not limited to leaf twisting may occur with some onion cultivars when Starane Ultra is applied as a broadcast treatment, especially when applications are made to larger dry bulb onions. Do not use Starane Ultra if the risk of injury is unacceptable.

### Specific Use Restrictions:

- Do not allow livestock to graze treated areas or harvest treated forage within 7 days of application.
- **Preharvest Interval (PHI):** Do not apply within 42 days of onion harvest.
- Do not make more than 2 applications per season.
- **Plant-Back Restriction:** Plant only labeled crops within 120 days of application.
- **Chemigation:** Do not apply through any type of irrigation system.
- Do not apply Starane Ultra when furrow irrigation is running. Manage treated fields to avoid water runoff for at least 6 hours after application.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow  
R357-034  
Accepted: 05/08/17  
Replaces: R357-008