

Supplemental Label

Armezon®

Herbicide

For use in sugarcane with associated rotational crop intervals

This supplemental label expires September 12, 2020, and must not be used or distributed after this date.

Active Ingredient:

topramezone: [3-(4,5-dihydro-isoxazolyl)-2-methyl-4-(methylsulfonyl)

phenyl](5-hydroxy-1-methyl-1*H*-pyrazol-4-yl)methanone 29.7%

Other Ingredients: 70.3%

Total: 100.0%

1 gallon contains 2.8 pounds of topramezone free acid.

EPA Reg. No. 7969-262

CAUTION/PRECAUCION

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Armezon® herbicide** container label, EPA Reg. No. 7969-262, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Armezon** before applying.
- Use of **Armezon** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Armezon**.

Product Information

Armezon provides systemic postemergence control or growth suppression of emerged broadleaf and grass weeds. Refer to the **Armezon** container label for a complete list of weeds controlled or suppressed and maximum weed sizes.

Sugarcane

Armezon can be applied to plant cane or sugarcane grown from stubble (ratoon). **Armezon** may be used in tank mixes or sequential applications with other herbicides registered for use in sugarcane such as atrazine, metribuzin, or **Prowl® H2O herbicide**. If **Armezon** is tank mixed with other herbicides, follow label restrictions for the most restrictive tank mix product. Application of **Armezon** may cause transient discoloration, chlorosis, or yellowing of sugarcane.

Armezon may be applied between growing seasons as either an early preplant in plant cane or post harvest in ratoon cane prior to cane initiating regrowth. Apply 0.5 to 2.0 fl ozs/A of **Armezon** with a minimum of 14 days between sequential applications. **DO NOT** apply more than 4.0 fl ozs/A of **Armezon** per year.

Special Weeds Controlled/Suppressed

In addition to the weeds controlled early postemergence as described in **Table 1** and **Table 2** of the container label, **Armezon** controls or suppresses the following weeds.

- **Common Bermudagrass** (*Cynodon dactylon*) – Apply early in the season at onset of Bermudagrass greenup or emergence of new leaves. Apply 1.0 to 2.0 fl ozs/A of **Armezon® herbicide** per application. Up to four sequential applications at 2 to 3 week intervals may be necessary for best control. **DO NOT** apply more than 4.0 fl ozs/A of **Armezon** (0.0875 lb topamezone per acre) per year. Apply using MSO or COC spray adjuvant plus nitrogen fertilizer such as AMS or UAN. See **Adjuvants** and **Nitrogen Fertilizer** sections for details.
- **Fall panicum** (*Panicum dichotomiflorum*) – For rescue suppression of large fall panicum more than 12-inches tall or other annual grasses listed in **Table 1** and **Table 2**, apply 2.0 fl ozs/A **Armezon** and use a minimum of 20 gallons per acre spray volume for proper spray coverage. Apply using MSO or COC spray adjuvant plus nitrogen fertilizer such as AMS or UAN. See **Adjuvants** and **Nitrogen Fertilizer** sections for details.

Sugarcane Restrictions

- **DO NOT** apply more than 2.0 fl ozs/A of **Armezon** (0.0438 lb topamezone/A) per application in sugarcane.
- **DO NOT** apply more than 4.0 fl ozs/A of **Armezon** (0.0875 lb topamezone/A) per year in sugarcane.
- **DO NOT** exceed a seasonal total of 2.0 fl ozs/A of **Armezon** during the final year of sugarcane production, prior to rotation to another crop.
- **DO NOT** apply **Armezon** within 100 days of sugarcane harvest.
- **DO NOT** graze or feed treated sugarcane for at least 100 days following an application of **Armezon**.

Rotational Crop Restrictions

The following rotational crops may be planted after applying **Armezon** at the specified rate. Planting earlier than the specified interval may result in crop injury. Avoid overapplication by minimizing overlap of spray swaths and by switching off spray boom when turning (end rows).

For rotational crops following the use of sequential applications of **Armezon**, the rotational interval begins after the last **Armezon** application.

Rotational Crop	Rotational Interval (months)		
	Armezon Application Rate (fl oz/A)		
	0.5	0.75	1.0 to 2.0
Corn (all)	0	0	0
Sugarcane	0	0	0
Cereal grains	3	3	3
Grass grown for seed	3	3	3
Rice	3	3	3
Alfalfa	9	9	9
Cotton	9	9	9
Grain sorghum	9	9	9
Peanut	9	9	9
Potato	9	9	9
Soybean	9	9	9
Sunflower	9	9	9
Canola	9	9	18
Lima bean, succulent	9	9	18
Dry beans ¹	9	9*	18**
Pea	9	9	18**
Snap/Garden bean	9	9*.†	18
Sugar beet	9*	9*	18**
All Other Crops	18	18	18

¹ For cranberry beans in Idaho, Utah, and the area east of the Cascade Mountains in Oregon and Washington, follow the guidelines for snap/garden bean.

* 18 months for the following states: Colorado, Michigan, Minnesota, Montana, Nebraska (west of Highway 83), North Dakota, South Dakota, Wisconsin, and Wyoming

** 9 months for Idaho, Oregon, and Washington

† 18 months for Idaho, Utah, and the area east of the Cascade Mountains in Oregon and Washington.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Armezon and Prowl are registered trademarks of BASF.

© 2017 BASF Corporation
All rights reserved.

007969-00262.20170316c.NVA 2017-04-216-0059

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry