

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 617A-S13 1016**

**Tilt®**

### Fungicide

**This supplemental label expires on October 21, 2019 and must not be used or distributed after this date.**

*Active Ingredient:*

Propiconazole*:	41.8%
-----------------	-------

Other Ingredients**:	58.2%
----------------------	-------

Total:	100.0%
--------	--------

\*CAS No. 60207-90-1

\*\*Contains petroleum distillates

Tilt formulation is an emulsifiable concentrate (EC)

Contains 3.6 lb of propiconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

**EPA Reg. No. 100-617**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Tilt as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## DIRECTIONS FOR USE

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Grasses grown for seed (Cool season grasses only)</b>	Rusts ( <i>Puccinia</i> spp.) Powdery Mildew ( <i>Erysiphe graminis</i> ) Ergot Stem Diseases	4-8	Apply Tilt when powdery mildew infections, and/or rust pustules are noticeable and increasing in number in late spring or early summer. Apply on a 14-21 day interval. To maximize control of severe rust pressure, apply 8 fl oz/A and make applications at 14-day intervals until the seed is mature. Make the last application at least 20 days before seed matures. For bluegrass, it is important to begin application early in the growing season.

**Application:** Tilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Tilt may be applied by ground, air (a minimum of 10 gal/A), or chemigation.

**Specific Use Restrictions:**

- 1) Do not feed hay cut within 20 days of the last application.
- 2) Do not graze treated areas within 140 days of the last application.
- 3) Do not apply more than 32 fl oz/A/season of Tilt.
- 4) Do not apply more than 0.90 lb ai propiconazole-containing products/A/season.
- 5) Do not apply to bermudagrass grown for seed.

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Mint Peppermint Spearmint	Rust ( <i>Puccinia menthae</i> )	4	Begin applications when the plants are 2-4 inches high or when conditions become favorable for disease development. Continue on a 14-day interval as conditions warrant.

**Application:** Tilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Tilt may be applied by ground in a minimum of 20 gal/A or by aerial in a minimum of 2 gal/A.

**Specific Use Restrictions:**

- 1) Do not apply within 7 days of harvest.
- 2) Do not exceed 12 fl oz of product/A/season.
- 3) Do not apply more than 0.338 lb ai propiconazole-containing product/A/season.

©2017 Syngenta

Tilt® and the Syngenta logo are trademarks of a Syngenta Group Company

Tilt 617A-S13 1016 – ep – 4-13-17