

# RESTRICTED USE PESTICIDE

DUE TO AVIAN AND AQUATIC TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

## DIAZINON AG500

GROUP

1B

INSECTICIDE

### Insecticide

For control of certain insects on fruits, nuts, vegetables, and ornamentals grown outdoors in nurseries

#### ACTIVE INGREDIENT:

% By Wt.

Diazinon: O,O-diethylO- (2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate . . . . . 48.0%

INERT INGREDIENTS\* . . . . . 52.0%

TOTAL . . . . . 100.0%

Contains 4 lbs. diazinon per gallon

\*Contains xylene range aromatic solvent

EPA Reg. No. 66222-9

EPA Est. No. 67545-AZ-001


KEEP OUT OF REACH OF CHILDREN  
CAUTION/  
PRECAUTION

For First Aid, Precautionary Statements, and Directions for Use, see inside of this booklet.

How can we help? 1-866-406-6262

Net Contents

2.5 gallon



INSECTICIDE

ADAMA

<b>FIRST AID</b> <b>Contains an organophosphate that inhibits cholinesterase</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>● Call a poison control center or doctor immediately for treatment advice.</li> <li>● Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>● Do not give any liquid to the person.</li> <li>● Do not give anything by mouth to an unconscious or convulsing person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>● Take off contaminated clothing.</li> <li>● Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>● Move person to fresh air.</li> <li>● If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>● Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>● Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Prosar at 1-877-250-9291 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN:</b> This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains petroleum distillates (xylene range aromatic solvent) which may pose an aspiration hazard. Vomiting may cause aspiration pneumonia. Gastric lavage may be indicated if product was taken internally.	

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contamination of food and feed. Food utensils such as tablespoons and measuring cups should not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Do not contaminate ornamental fish ponds.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below.

**Mixers, loaders, applicators, and other handlers using engineering controls must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of barrier laminate or Viton ≥14 mils if mixing or loading
- Chemical-resistant apron if mixing or loading

See engineering controls for additional requirements.

**Handlers performing tasks, such as cleaning equipment or spill cleanup for which engineering controls are not feasible must wear:**

- Coveralls over long-sleeved shirt and long pants
- Chemical resistant gloves made of barrier laminate or Viton ≥14 mils
- Chemical-resistant footwear plus socks
- Chemical-resistant apron if exposed to the concentrate
- Goggles, face shield, or safety glasses
- Powered air purifying canister-type respirator (gas-mask) equipped with an organic vapor canister that incorporates HE filters (NIOSH approval number prefix TC-14G).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

#### ENGINEERING CONTROLS

Mixers and loaders must use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides that provides dermal and inhalation protection [40 CFR 170.240 (d)(4)] and must:

--wear the personal protective equipment required for mixers/loaders using engineering controls;

--wear protective eyewear if the system operates under pressure; and

--be provided and have immediately available for use in an emergency such as a broken package, spill, or equipment breakdown, chemical-resistant footwear and the respirator specified in the PPE section of this labeling for handlers not using engineering controls.

Applicators using motorized ground equipment must use an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural

Pesticides [40 CFR 170.240 (d) (5)] for dermal protection. Flaggers supporting aerial applications to lettuce must use an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240 (d) (5)] for dermal protection.

In addition, applicators must:

- wear the personal protective equipment required in the PPE section of this labeling for handlers using engineering controls;
- either wear the respirator specified for handlers not using engineering controls or use an enclosed cab that is declared in writing by the manufacturer or by a government agency to provide at least as much respiratory protection as the respirator specified for handlers not using engineering controls;
- be provided and have immediately available for use in an emergency when they must exit the cab in the treated area: coveralls, chemical-resistant footwear, and, if not already using one, the respirator specified for handlers not using engineering controls;
- take off any PPE that was worn in the treated area before reentering the cab, and
- store all such PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

**Exception:** For application to lettuce, see directions for use for a special exception to these engineering controls requirements.

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (6)].

When entering or leaving an aircraft contaminated with pesticide residues, pilots must wear chemical-resistant gloves and must store used gloves in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cockpit.

**Note:** Aerial applications are permitted only on lettuce.

#### USER SAFETY RECOMMENDATIONS

##### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwater. This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

#### PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use, pour, spill, or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Reformulation or repackaging of this product is prohibited.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI). The REI for each crop is listed in the directions for use associated with each crop. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard under certain circumstances allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls worn over long sleeve shirt and long pants
- Chemical resistant gloves made of barrier laminate or Viton  $\geq 14$  mils
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear if overhead exposures

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

## WORK SAFETY RULES

Repeated exposures to cholinesterase inhibitors such as are contained in this product may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. When handling Diazinon AG500, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See **FIRST AID** section of this label. When handling this product, wear the personal protective equipment listed on this label.

## GENERAL INFORMATION

Diazinon AG500 is an emulsifiable solution which, when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops and other plants. Do not use in greenhouses. Do not use in barns or dwellings. Do not use on livestock. Aerial application is prohibited, except as allowed on the label for use on lettuce.

In the directions for use which follow, best control is obtained when application is made when infestation is noticed.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

Do not use water from treated cranberry beds for drinking purposes or to irrigate crops other than those appearing on EPA approved diazinon labels.

**Note:** The California Department of Pesticide Regulation has set a 5-day reentry period for diazinon on peaches and nectarines. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

Failure to follow the directions for use and precautions on this label may result in poor insect control, crop injury, or illegal residues.

## SPRAY EQUIPMENT

Applications of Diazinon AG500 using sufficient water volume to provide thorough and uniform coverage generally provide the most effective pest control.

To avoid spray drift, do not apply when conditions favor drift from beyond the target area. Avoid overlap because crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate sprayer before use.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult your local or state agricultural authorities.

## MIXING PROCEDURES

Thoroughly clean spray equipment before using this product. Prepare only the amount of spray mixture needed for the immediate application. Vigorous agitation is necessary for proper dispersal of this product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand in the spray tank overnight. Flush the spray equipment thoroughly following each use.

### 1. Diazinon AG500 Alone:

Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the Diazinon AG500 to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the Diazinon AG500 has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

### 2. Diazinon AG500 in Tank Mixtures:

**Note: Do not mix Diazinon AG500 with any formulation of captan or Captec® because crop injury may occur.** Diazinon is physically compatible with most insecticide and fungicide products; if you are not certain of the physical compatibility of Diazinon AG500 with other tank mix partners, please contact your local ADAMA sales representative. When using Diazinon AG500 in tank mixtures, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix products' labels. Confirm that the tank mixture is safe to the crop by spraying a small area first and evaluating crop safety after an appropriate period. Do not treat larger areas until crop safety has been confirmed.

Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add tank mix partners in this order: all products in water-soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables, liquids, and emulsifiable concentrates. Always allow each tank mix partner to fully disperse before adding the next product. Maintain sufficient agitation while adding the remainder of the water and until all of the mixture has been applied.

## CHEMIGATION STATEMENT

### APPLICATION THROUGH IRRIGATION SYSTEMS - CHEMIGATION

Diazinon AG500 may be applied alone or in combination with other pesticides registered for application through irrigation systems. Apply this product only through sprinkler including center pivot, lateral move, end tow side (wheel) roll, traveler, big gun, solid set, or hand move, flood (basin) furrow, border, or drip (including surface and subsurface) irrigation systems. Do not apply this product through any other type of irrigation system.

Diazinon is physically compatible with most insecticide and fungicide products; if you are not certain of the physical compatibility of Diazinon AG500 with other tank mix partners, please contact your local ADAMA sales representative. **Note: Do not mix Diazinon AG500 with any formulation of captan or Captec® because crop injury may occur.**

Crop injury, lack of effectiveness, or illegal other pesticides, surfactants, or fertilizers unless prior use has shown the combination noninjurious under your conditions of use. Follow precautionary statements and directions for all tank mix products.

On all crops, use sufficient gallage of water to obtain thorough and uniform coverage but not cause runoff or excessive leaching. This will vary depending on equipment, pest problem, and stage of crop growth. Application of more or less than optimal quantity of water may result in decreased chemical performance, crop injury, or illegal pesticide residues.

Meter this product into the irrigation water uniformly during the period of operation. Do not overlap application. Follow recommended label rates, application timing, and other directions and precautions for crop being treated. Continuous mild agitation of pesticide mixture may be needed to assure a uniform application, particularly if the supply tank requires a number of hours to empty.

Do not apply when wind speed favors drift beyond the area intended for treatment.

## CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

*Note: ADAMA does not encourage connecting chemigation systems to public water supplies. The following information is provided for users who have diligently considered all other application and water supply options before electing to make such a connection.*

Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ), or the functional equivalent in the water supply line upstream from a point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

### SPRINKLER CHEMIGATION

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

### OPERATING INSTRUCTIONS

1. Determine the acreage to be treated by the sprinkler system.
2. Measure the appropriate amount of Diazinon AG500 needed to treat the acreage to be treated. Refer to the specific directions for use for the application rate and the amount of water per acre.
3. Add the premeasured Diazinon AG500 to the chemigation system supply tank and dilute with water if needed. Proper agitation is needed if Diazinon AG500 is tank mixed with other pesticides registered for sprinkler chemigation application.
4. Start the irrigation system and bring the system up to full pressure making certain that all sprinkler heads are functioning properly.
5. Engage the chemigation injection or venturi system to add the diazinon to the irrigation lines.
6. Continue to run irrigation system for a few minutes after supply tank is empty to allow material in the irrigation lines to reach the sprinkler heads furthest from the chemigation injection point.

### FLOOD (BASIN) FURROW AND BORDER CHEMIGATION (SOIL DRENCH USES)

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- a) The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- b) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- c) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- d) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- e) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- f) Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

### DRIP (INCLUDING SURFACE AND SUBSURFACE) CHEMIGATION (SOIL DRENCH USES)

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pipe and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

#### PREHARVEST INTERVAL

The required days to wait between the last application and harvest are given in ( ) after each crop name.

#### FRUIT AND NUT CROPS

Diazinon AG500 may be applied using ground application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for application. The rate of Diazinon AG500 for concentrated spray applied to fruit trees by ground equipment is based on a dilute full cover spray applied with conventional ground equipment at a rate of 300-400 gals. of water per acre. The specific amount of product to be used appears under the separate directions for use for each crop. If the rate calls for 1 pt. of Diazinon AG500 in 100 gals. of water, this amounts to 3-4 pts. of product per acre when applied at the usual dilute ground spray volumes of 300-400 gals. of water per acre. In no case should the amount of product used per acre exceed the maximum amount per acre per application that is specified in the **COMMENTS** section for individual crops.

**Note:** To protect bees, do not apply this product to fruit trees when trees or substantial numbers of weeds in the orchard are in bloom.

Diazinon AG500 may be used during dormancy on deciduous trees and vines. When mixing with spray oils, follow oil manufacturer's use directions.

Crop	Ground Application Minimum Gals./Acre	
	Dilute	Concentrate
Almonds*	100	20
Apricots	100	20
Blackberries**	100	20
Blueberries	100	20
Boysenberries**	100	20
Caneberries****	100	20
Cherries	100	20
Cranberries	15	-
Dewberries**	100	20
Figs*	100	20
Filberts***	250	20
Loganberries**	100	20
Nectarines	100	20
Peaches	100	20
Plums	100	20
Prunes	100	20
Raspberries**	100	20
Strawberries	100	20

\*CA Only

\*\*CA, OH, OR, WA Only

\*\*\*OR, WA Only

\*\*\*\* CA, OH, OR, WA Only

**FOR PROTECTION OF AQUATIC SPECIES IN COUNTIES AND PORTIONS OF COUNTIES LOCATED IN THE SACRAMENTO AND SAN JOAQUIN VALLEYS BELOW 1000 FEET ELEVATION FROM POTENTIAL CONTAMINATION IN SURFACE WATERS OF DIAZINON APPLIED DURING THE DORMANT SPRAY PERIOD, THE FOLLOWING ADDITIONAL LABEL REQUIREMENTS APPLY:**

- Dormant applications on orchard crops are restricted to ground application equipment only.
- Do not apply within 100 feet upslope of "sensitive aquatic sites" such as any irrigation ditch, drainage canal, or body of water that may drain into a river or tributary unless a suitable method is used to contain or divert runoff waters. Waters that are contained or diverted must be held for a minimum of 72 hours before release into a sensitive aquatic site.
- Maintain a vegetative buffer strip a minimum of 10 feet wide from the edge of a field that is adjacent to and within 100 feet of sensitive aquatic sites.
- Do not apply this product to orchards when soil moisture is at field capacity and/or when a storm event likely to produce runoff from the treated orchard is forecasted by NOAA/NWS (National Weather Service) to occur within 48 hours following application.
- Make dormant applications only when insect scouting information or the recommendation of a Pest Control Advisor indicates treatment is required. (See UC IPM Guidelines for San Jose Scale in stone fruits and almonds and aphids in stone fruits. Use the prune dormant spur sampling program to determine need for a dormant treatment in that crop).

- Apply only when wind speed is 3-10 mph at the application site as measured by an anemometer outside of the orchard on the side nearest and upwind from a sensitive site.
- When sensitive aquatic sites are downwind from orchards, spray the first three rows nearest the sensitive aquatic sites only when the wind is blowing away from the sites. The row at the edge of the field next to sensitive aquatic sites must be sprayed with the outside nozzles turned off. Spray must not be directed higher than the tree canopy, and spray must be directed away from sensitive aquatic sites.
- The Stewardship Bulletin **“Orchard Practices for Protecting Surface Water”** must be available to handlers and equipment operators at the application site during all application activities. (Note: Bulletin available through CURES.)

#### FRUIT AND NUT CROPS

CROP	PEST	RATE	COMMENTS
<b>ALMONDS</b> (California Only)	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borers, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	1-1 ½ pts. per 100 gals. water	Apply in 100 gals. water plus 2-3 gals. dormant oil or 1-1½ gals. superior type oil. Apply during the dormant season only.
	<ul style="list-style-type: none"> <li>• Do not apply more than 6 pts. of product (3 lb a.i.) or 9 gals. of oil per acre.</li> <li>• Do not apply more than 3 lbs. of a.i. per acre per year.</li> <li>• Make a maximum of one application per year during the dormant season only.</li> <li>• Apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>• The REI is 7 days.</li> </ul>		
<b>APRICOTS</b>	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	1 pt. per 100 gals. water	Apply as a dormant spray only once per year. When using spray oils, follow spray oil manufacturer's use directions.
	Aphids, Brown Mite (Clover Mite), Two-spotted Spider Mite, Olive Scale Crawler, San Jose Scale Crawler	1 pt. per 100 gals. water	Apply when infestation is noticed.
	Olive Scale Crawler	1/2 pt. plus 1½ gals. of light medium horticultural oil per 100 gals. of water	Apply when crawlers are present.
	Apricot Mealybug	1 pt. per 100 gals. water	Apply as a cover spray from petal fall to June.
	<ul style="list-style-type: none"> <li>• Do not apply more than 4 pts. of product (2 lb a.i.) per acre per application.</li> <li>• Do not apply more than 4 lbs. of a.i. per acre per year.</li> <li>• A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>• For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>• The REI is 4 days.</li> <li>• The PHI is 21 days.</li> <li>• Allow 60 days between dormant application and in-season application; 120 days between dormant application and post-harvest application</li> </ul>		
<b>BLUEBERRIES</b>	Cranberry Fruitworm, Cherry Fruitworm, Blueberry Maggot, Aphids, Thrips, Two-spotted Spider Mite	1 pt. per 100 gals. water	Apply when infestation occurs.
	Fire Ants	1 pt. per 100 gals. water	To aid in the control of fire ants, slowly apply 1 gal. of diluted mixture over and 6 inches around each mound. Apply gently to avoid disturbing ants. Use equipment capable of delivering the diluted product as a gentle rain. High pressure sprays may disturb the ants and cause migration reducing product effectiveness. For best results, apply in cool weather (65-80°F) or in early morning or late evening hours.
	<ul style="list-style-type: none"> <li>• Do not apply more than 2 pts. of product (1 lb a.i.) per acre per application.</li> <li>• Do not apply more than 2 lbs. of a.i. per acre per year.</li> <li>• A maximum of two applications are allowed per year: 1) a maximum of one as an in-season foliar application regardless of target pest and 2) a maximum of one for control of fire ant.</li> <li>• The REI is 5 days.</li> <li>• The PHI is 7 days.</li> <li>• Allow a minimum of 30 days between treatments.</li> </ul>		

CROP	PEST	RATE	COMMENTS
<b>CANEBERRIES</b> (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries) (California, Ohio, Oregon, and Washington Only)	Raspberry Fruitworm	1 qt. per acre	Apply when insects occur. Apply in a minimum of 100 gals. of water per acre (200 gals. per acre maximum). Make application when blossom buds separate.
	Raspberry Crown Borer (Raspberry Root Borer)	2 qts. per acre	Apply a single application in a minimum of 100 gals. of water per acre as a drench to the crown and lower canes in the spring before buds break.
	<ul style="list-style-type: none"> <li>Do not apply more than 2 qts. of product (2 lb a.i.) per acre per application.</li> <li>Do not apply more than 2 lbs. of a.i. per acre per year.</li> <li>Make a maximum of one application per year regardless of the target pest.</li> <li>The REI is 5 days.</li> <li>The PHI is 7 days.</li> </ul>		
<b>CHERRIES</b>	San Jose Scale, Parlortaria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	1 pt. per 100 gals. water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Apply only once during the dormant season.
	Eyespotted Bud Moth, Fruitree Leafroller, Leafhoppers, Black Cherry Aphid, San Jose Scale Crawler	1 pt. per 100 gals. water	Apply when infestation occurs.
	Cherry Fruit Fly	1/2 - 1 pt. per 100 gals. water	Apply when adult flies begin to emerge.
	Cherry Rust Mite	1/2 - 1 pt. per 100 gals. water	Apply to trees after harvest.
	<ul style="list-style-type: none"> <li>Do not apply more than 4 pts. of product (2 lb. a.i.) per acre per application.</li> <li>A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>Do not apply more than 8 pints (4 lbs. of a.i.) per acre per year</li> <li>For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>The REI is 4 days.</li> <li>The PHI is 21 days.</li> <li>Allow 30 days between dormant application and in-season application; 90 days between dormant application and post-harvest application.</li> </ul>		
<b>CRANBERRIES</b>	Blackheaded Fireworm ( <i>Rhopobota naevana</i> )	2 qts. per acre	Apply to larval stage. Pheromone trap captures and sweep monitoring may be used to optimize treatment timing. Repeat application after 7 days if necessary. A maximum of 3 applications per season is permitted regardless of target pest.
	Cranberry Fruitworm	2 to 3 qts. per acre	Cranberry growth stage and/or berry inspection for eggs may be used to optimize treatment timing; consult your local extension agent or pest management adviser. Repeat application after 7 days if necessary. A maximum of 3 applications per season is permitted regardless of target pest.
	Cranberry Tipworm	2 qts. per acre	Presence of damage is not an indicator of active infestation. For best results, target early-stage larvae. Repeat application after 14 days if necessary. A maximum of 3 applications per season is permitted regardless of target pest.
	<ul style="list-style-type: none"> <li>Do not make more than 3 applications per year regardless of the target pest.</li> <li>Do not use water from irrigated or flooded cranberry bed for drinking purposes or to irrigate crops other than those appearing on EPA-approved diazinon labels.</li> <li>Do not apply more than 3 qts. (3 lbs.) of this product per acre per application.</li> <li>Do not apply more than 18 pints of this product (9 lbs. a.i.) per acre per year.</li> <li>Allow a minimum of 14 days between treatments</li> <li>The REI is 5 days.</li> <li>The PHI is 7 days.</li> </ul>		



CROP	PEST	RATE	COMMENTS
<b>FIGS</b> (California Only)	Vinegar flies- <i>Drosophila</i> spp., Dried fruit beetle	1 pt. per 100 gals.	Apply 1 pt. per 100 gals. water not to exceed 100 gals. of spray per acre. Apply when infestation occurs.
	<ul style="list-style-type: none"> <li>Do not apply more than 1 pt. of product (0.5 lb a.i.) per acre.</li> <li>Make a maximum of one application per year regardless of target pest.</li> <li>Do not apply more than 0.5 lbs. of a.i per acre per year.</li> <li>Maximum yearly application is 0.5 lbs.</li> <li>The REI is 4 days.</li> <li>The PHI is 21 days.</li> </ul>		
<b>FILBERTS</b> <b>(HAZELNUTS)</b> (Oregon and Washington Only)	Filbert Leafroller, Aphids	1 pt.	Apply as a thorough cover spray in 250 to 400 gals. of water per acre.
	<ul style="list-style-type: none"> <li>Do not apply more than 1 pint of product (0.5 lb a.i.) per acre.</li> <li>Make a maximum of one application per year regardless of target pest.</li> <li>The REI is 18 days.</li> <li>The PHI is 45 days.</li> <li>Do not apply more than 0.5 lbs. of a.i per acre per year.</li> </ul>		
<b>NECTARINES</b>	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	1 pt. per 100 gals. water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Apply only once during the dormant season.
	Oriental Fruit Moth, Aphids, Brown Mite (Clover Mite), Two-spotted Spider Mite, Olive Scale Crawler, San Jose Scale Crawler, Peach Twig Borer	1 pt. per 100 gals. water	Apply when infestation occurs. Do not apply more than 4 pts. of product or 4 gals. of summer oil per acre.
	<ul style="list-style-type: none"> <li>Do not apply more than 4 pts. of product (2 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs. of a.i. per acre per year.</li> <li>A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>The REI is 4 days.</li> <li>The PHI is 21 days.</li> <li>Allow 60 days between dormant application and in-season application.</li> </ul>		
<b>PEACHES</b>	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	1 pt. per 100 gals. water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Do not apply more than 4 pts. of product or 6 gals. of oil per acre. Apply only once during the dormant season.
	Brown Mite (Clover Mite), Two-spotted Spider Mite, Olive Scale Crawler, San Jose Scale Crawler	1 pt. per 100 gals. water	Apply when infestation occurs.
	Peach Twig Borer, Parlatoria Scale	1 pt. per 100 gals. water	Apply when infestation occurs.
	Oriental Fruit Moth	1 pt. per 100 gals. water	Apply at 100% petal fall.
	Leafhoppers	1 pt. per 100 gals. water	Apply when infestation occurs.
	White Peach Scale	1-1 1/3 pts./100 gals. water	Apply post harvest timed to coincide with peak crawler and immature scale activity. Do not apply before fruit is harvested.
	<ul style="list-style-type: none"> <li>Do not apply more than 4 pts. of product (2 lb a.i.) per acre per application.</li> <li>A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>Do not apply more than 4 lbs. of a.i per acre per year.</li> <li>The REI is 4 days.</li> <li>The PHI is 21 days.</li> <li>Allow 60 days between dormant application and in-season application; 120 days between dormant application and post-harvest application.</li> </ul>		

CROP	PEST	RATE	COMMENTS
PLUMS	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	1 pt. per 100 gals. water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Apply only once during the dormant season.
	Brown Mite (Clover Mite), European Red Mite	1 pt. per 100 gals. water	Apply when infestation occurs.
	Leafcurl Plum Aphid, Mealy Plum Aphid, Thistle Aphid	1/2 to 1 pt. per 100 gallons water	Apply when infestation occurs
	<ul style="list-style-type: none"> <li>Do not apply more than 4 pts. of product (2 lb a.i.) per acre per application.</li> <li>A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>Do not apply more than 4 lbs. of a.i. per acre per year.</li> <li>For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>The REI is 4 days.</li> <li>The PHI is 21 days.</li> <li>Allow 60 days between dormant application and in-season application; 120 days between dormant application and post-harvest application.</li> </ul>		
PRUNES	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	1 pt. per 100 gals. water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Apply only once during the dormant season.
	Brown Mite (Clover Mite), European Red Mite	1 pt. per 100 gals. water	Apply when infestation occurs.
	Leafcurl Plum Aphid, Mealy Plum Aphid, Thistle Aphid	1/2 - 1 pt. per 100 gals. water	Apply when infestation occurs.
	<ul style="list-style-type: none"> <li>Do not apply more than 4 pts. of product (2 lb a.i.) per acre per application.</li> <li>A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>Do not apply more than 4 lbs. of a.i. per acre per year.</li> <li>For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>The REI is 4 days.</li> <li>The PHI is 21 days.</li> <li>Allow 60 days between dormant application and in-season application; 120 days between dormant application and post-harvest application.</li> </ul>		
STRAWBERRIES	Aphids, Two-spotted Spider Mite,	1 pt. (0.5 lb a.i.) per 100 gals. water per acre	Apply when pest occurs.
	Cyclamen Mite	1 qt. (1 lb a.i.) per 100 gals. water per acre	Direct spray to the plant crown. Plants should be sufficiently agitated to assure thorough coverage of the foliage.
	Mole Crickets	1 qt. (1 lb a.i.) per acre	Broadcast in sufficient water to obtain even coverage when insects are present. Apply 1-2 days before transplanting and immediately incorporate into the top 1-2 inches of soil.
	Strawberry Leafroller	0.75 - 1 pint per 100 gals water	Apply 200 gallons of spray per acre when blossoms show color
	<ul style="list-style-type: none"> <li>Do not apply more than 1 qt. of product (1 lb a.i.) per acre per application.</li> <li>Make a maximum of one foliar application per crop and a maximum of one soil application per crop.</li> <li>Do not apply more than 2 lbs. of a.i. per acre per year.</li> <li>Allow a minimum of 30 days between treatments.</li> <li>The REI is 3 days.</li> <li>The PHI is 5 days.</li> </ul>		

### VEGETABLE CROPS

For foliar or soil applications, Diazinon AG500 should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. Diazinon AG500 may be applied using ground or aerial (lettuce only) application equipment as specified in the following table. Aerial application is prohibited, except as allowed on the label for use on lettuce. This table indicates the minimum amount of water that can be used for application. The specific amount of product to be used appears under the separate directions for use for each crop.

CROP	Ground Application Min. GPA	Min. Gals./1000 Sq. Ft. (Ground)	AERIAL APPLICATION MIN. GPA
Beans, Succulent	10	2	-
Beets, Red (Table)	10	2	-
Broccoli	10	2	-
Brussels Sprouts	10	2	-
Cabbage	10	2	-
Cantaloupes	5	1	-
Carrots	10	2	-
Casaba Melons	5	1	-
Cauliflower	10	2	-
Collards	10	2	-
Crenshaw Melons	5	1	-
Endive (Escarole)	10	2	-
Ginseng	10	2	-
Honeydew Melons	5	1	-
Kale	10	2	-
Lettuce, Head	10	2	5
Lettuce, Leaf	10	2	5
Muskmelons	5	1	-
Mustard Greens	10	2	-
Onions (Bulb and Green)	10	2	-
Peas, Succulent	10	2	-
Persian Melons	5	1	-
Radishes	10	2	-
Rutabagas	10	2	-
Spinach	10	2	-
Tomatoes	10	2	-
Watermelons	5	1	-

Observe stated time intervals between last application and harvest as well as intervals between and maximum numbers of applications per season. Do not apply in greenhouses.

NOTES: (1) Soil Incorporation: Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see directions for individual crops) using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches; for subterranean cutworms, incorporate 3-6 inches. (2) Diazinon will not control organophosphate-resistant leafminers.

### VEGETABLE CROPS

CROP	PEST	RATE/ACRE	COMMENTS
<b>BEANS, SUCCULENT</b>	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lb of a.i. per acre per year</li> <li>The REI is 3 days.</li> <li>The PHI is 45 days.</li> </ul>		

CROP	PEST	RATE/ACRE	COMMENTS
<b>BEEETS, RED</b> (Table)	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 3 days.</li> </ul>		
<b>BROCCOLI, BROCCOLINI</b>	Root Maggots	2-3 qts.	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. per 50 gals. water	Apply in transplant water as a drench application when 200-300 gals of water are used per acre. Apply recommended rate by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i. per acre per year.</li> <li>The REI is 4 days.</li> </ul>		
<b>BRUSSELS SPROUTS</b>	Root Maggots	2-3 qts.	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. per 50 gals. water	Apply in transplant water as a drench application when 200-300 gals of water are used per acre. Apply recommended rate by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i. per acre per year.</li> <li>The REI is 4 days.</li> </ul>		
<b>CABBAGE</b>	Root Maggots	2 - 3 qts.	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. per 50 gals. water	Apply in transplant water as a drench application when 200-300 gals of water are used per acre. Apply recommended rate by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1 - 2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 4 days.</li> </ul>			

CROP	PEST	RATE/ACRE	COMMENTS
CARROTS	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 3 days.</li> </ul>
CAULIFLOWER, BROCCOFLOWER	Root Maggots	2-3 qts.	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		4-8 oz. per 50 gals. water	Apply in transplant water as a drench application when 200-300 gals of water are used per acre. Apply recommended rate by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 4 days.</li> </ul>
COLLARDS	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 4 days.</li> </ul>
ENDIVE (Escarole)	Cutworms	1 qt.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	1 qt.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 2pts. of product (1lb a.i.) per acre per application.</li> <li>Do not apply more than 2 pts (1lb a.i) a.i per acre per year.</li> <li>The REI is 4 days.</li> </ul>
GINSENG	Leafhoppers, Aphids, Lygus Bugs, Flea Beetles, Jumping Plant Lice	3/4-1 pt.	Spray when insects first appear. For heavy insect infestations, use 1 pt. per acre. Apply by ground equipment using 10-100 gals. of water per acre.
			<ul style="list-style-type: none"> <li>Do not apply more than one application per year.</li> <li>Do not apply more than 1 pt. of product (0.5 lb a.i.) per acre per application</li> <li>Do not apply more than 0.5 lbs a.i. per acre per year.</li> <li>Do not apply during flowering of three- or four-year-old crops.</li> <li>Do not graze treated areas or feed treated forage to livestock.</li> <li>The REI is 3 days.</li> <li>The PHI is 30 days.</li> </ul>

CROP	PEST	RATE/ACRE	COMMENTS
<b>KALE</b>	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 4 days.</li> </ul>
<b>LETTUCE</b> (Head and Leaf)	Cutworms	2 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	2 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminer	1/2 - 1 pt.	Apply when insects occur.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per crop regardless of target pest.</li> <li>Do not make more than one foliar application per crop regardless of target pest.</li> <li>Do not apply more than 2 qts. of product (2 lb a.i.) per acre per application.</li> <li>Allow at least 30 days between applications.</li> <li>Aerial application is permitted on lettuce.</li> <li>EXCEPTION: Applicators applying to lettuce are permitted to use open-cab equipment provided they are wearing chemical-resistant headgear AND the personal protective equipment specified in this labeling for handlers not using engineering controls.</li> <li>The REI is 3 days.</li> <li>The PHI is 14 days.</li> <li>Do not apply more than 4 lbs a.i. per acre per year.</li> </ul>
<b>MELONS</b> (Cantaloupes, Casabas, Crenshaws, Muskmelons, Persians and hybrids of these) and <b>WATERMELONS (3)</b>	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 8 pints of this product (4 lbs. a.i.) per acre per year.</li> <li>The REI is 3 days.</li> <li>The PHI is 3 days.</li> </ul>
<b>HONEYDEW MELONS</b>	Aphids, Dipterous Leafminer, Thrips, Striped Cucumber Beetle, Melonworms, Leafhoppers, Spider Mites	1/2-1½ pts.	Apply when infestation occurs.
			<ul style="list-style-type: none"> <li>Do not make more than 1 soil and 1 foliar application per year regardless of target pest.</li> <li>Minimum retreatment interval is 30 days.</li> <li>For foliar application, do not apply more than 1.6 pints of this product (0.8 lb. a.i.) per acre per application.</li> <li>Do not apply more than 1.6 pints of this product (0.8 lb a.i) per acre per year.</li> <li>The REI is 3 days.</li> <li>The PHI is 3 days.</li> </ul>
<b>MUSTARD GREENS</b>	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.

CROP	PEST	RATE/ACRE	COMMENTS
<b>MUSTARD GREENS</b>	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 8 pints of this product (4 lbs. a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 4 days.</li> </ul>
<b>ONIONS</b> (Bulb and Green), <b>GARLIC, LEEKS,</b> <b>SPRING ONIONS</b> <b>OR SCALLIONS,</b> <b>JAPANESE</b> <b>BUNCHING</b> <b>ONIONS, GREEN</b> <b>SHALLOTS, GREEN</b> <b>ESCHALOTS</b>	Onion Maggots	2-4 qts.	Broadcast just before planting and mix into the top 3-4 inches of soil. Note: Diazinon will not control organophosphate-resistant onion maggots.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 8 pts. of product (4 lb a.i.) per acre per application.</li> <li>The REI is 3 days.</li> <li>Do not apply more than 4 lbs. a.i. per acre per year</li> </ul>
<b>RADISHES</b>	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 3 days.</li> </ul>
<b>RUTABAGAS</b>	Cabbage Maggot	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil to a depth of 4 inches.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application.</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 4 days.</li> </ul>
<b>SPINACH</b>	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application</li> <li>Do not apply more than 4 lbs a.i per acre per year.</li> <li>The REI is 3 days.</li> <li>The MRI is 1 day.</li> </ul>
<b>TOMATOES</b>	Cutworms	2-4 qts.	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	1 qt.	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
			<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>Do not apply more than 4 qts. of product (4 lb a.i.) per acre per application</li> <li>Do not apply more than 8 pints of this product (4 lbs. a.i.) per acre per year.</li> <li>The REI is 2 days.</li> </ul>

#### ORNAMENTAL STATEMENT

Before treating a large number of ornamental plants with Diazinon AG500 alone or as a tank mixture with any other material, make a test application on a few plants and observe for 7 to 10 days prior to treating large areas to reduce the possibility of plant injury.

## INSECT CONTROL ON ORNAMENTALS GROWN OUTDOORS IN NURSERIES

CROP	PEST	RATE	COMMENTS
<b>ARBORVITAE, AZALEA, BALSAM FIR (MAINE ONLY), BIRCH, BOXWOOD, CAMELLIA, CARNATION, CHRYSANTHEMUM, CRABAPPLE, DOGWOOD, DOUGLAS FIR, ELM, EUONYMUS, GLADIOLI, HAWTHORN, HOLLY, HONEY LOCUST, HONEYSUCKLE, JUNIPER, LILAC, LILY, LOCUST, MAPLE, MARIGOLD, OAK, PANSY, PETUNIA, PINE, ORNAMENTAL PLUM, POPLAR, RHODODENDRON, ROSE, SCOTCH PINE, SPIREA, SPRUCE, SYCAMORE, WAX PLANT (Hoya), WILLOW, YELLOWWOOD, and YEW</b>	Aphids, Bagworms, Carnation Bud Mite, Carnation Shoot Mite, Clover Mite, Cyclamen Mite, Dipterous Leafminer, European Pine Shoot Moth, European Red Mite, Flea Beetles, Holly Bud Moth, Leafhoppers, Obscure Root Weevil, Omnivorous Leafminer, Privet Mite, Scale Crawlers (Cottony-cushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Two-spotted Spider Mite, Juniper Webworm, Whiteflies	1/2 fl. oz. per 3 gals. water or 1 pt. per 100 gals. of water	Apply the recommended rates. Spray when listed pests first appear. Try to spray underside of leaves and penetrate dense foliage.
	Apple-and-Thorn Skeletonizer, Cotonest Webworm, European Pine Sawfly, Fall Webworm, Hemlock Cherries, Oak Looper, Obliquebanded Leafroller, Pear Slug, Tent Caterpillar	1½ fl. oz. per 3 gals. of water or 2 pts. per 100 gals. of water	
	Mimosa Webworm	1/2 fl. oz. in 3 gals of water or 1 pt. in 100 gals. of water	Apply as a thorough foliar spray when adults first appear. Try to spray underside of leaves and penetrate dense foliage.
<ul style="list-style-type: none"> <li>Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, and Gardenia since injury to the plants may occur.</li> <li><b>Note:</b> 1 fl. oz. = 2 tablespoons; 16 fl. oz. = 1 pint</li> <li>The maximum application rate on all ornamentals is 2 pts. (1 lb a.i.) per acre.</li> <li>Make a maximum of one application per crop cycle.</li> <li>The REI is 7 days for flowers and other commercial ornamentals grown for cutting.</li> <li>The REI is 2 days for all other commercial ornamentals.</li> <li>Application is permitted only on commercial ornamentals grown outdoors in nurseries.</li> <li>Do not apply more than 1 lb a.i. per crop cycle.</li> </ul>			

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a dry place. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. To avoid harming aquatic organisms in rivers and other surface waters, do not pour the concentrate, spray mixture, or rinse water into sanitary drains (for example, street drains). If pesticide, spray mixture, or rinsate cannot be disposed of by use according to the label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### CONTAINER DISPOSAL:

**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill, incinerate or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

**Nonrefillable Container (greater than five gallons):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill, incinerate or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

### LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **Conditions, Disclaimer of Warranties and Limitations of Liability.**

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ADAMA Makhteshim, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ADAMA Makhteshim, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of ADAMA Makhteshim, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ADAMA disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at ADAMA Makhteshim, Inc.'s election, the replacement of product.

**Captec 4L®** is a registered trademark of Arysta LifeScience North America LLC

042017v2



# RESTRICTED USE PESTICIDE

## DUE TO AVIAN AND AQUATIC TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

# DIAZINON AG500

### Insecticide

For control of certain insects on fruits, nuts, vegetables, and ornamentals grown outdoors in nurseries

### ACTIVE INGREDIENT:

% By Wt.

Diazinon: O,O-diethylO-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate . . . . . 48.0%

INERT INGREDIENTS\* . . . . . 52.0%

TOTAL . . . . . 100.0%

Contains 4 lbs. diazinon per gallon

\*Contains xylene range aromatic solvent

EPA Reg. No. 66222-9

EPA Est. No. 67545-AZ-001

## KEEP OUT OF REACH OF CHILDREN CAUTION/ PRECAUTION

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contamination of food and feed. Food utensils such as tablespoons and measuring cups should not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Do not contaminate ornamental fish ponds.

**How can we help?**  
**1-866-406-6262**

**Manufactured for:**  
**Makhteshim Agan of North America, Inc.**  
**(d/b/a ADAMA)**  
**3120 Highwoods Blvd.**  
**Raleigh, NC 27604**

Net Contents  
**2.5 gallons**



INSECTICIDE

**ADAMA**

GROUP

1B

INSECTICIDE

### FIRST AID

#### Contains an organophosphate that inhibits cholinesterase

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious or convulsing person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Prosar at 1-877-250-9291 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains petroleum distillates (xylene range aromatic solvent) which may pose an aspiration hazard. Vomiting may cause aspiration pneumonia. Gastric lavage may be indicated if product was taken internally.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a dry place. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. To avoid harming aquatic organisms in rivers and other surface waters, do not pour the concentrate, spray mixture, or rinse water into sanitary drains (for example, street drains). If pesticide, spray mixture, or rinsate cannot be disposed of by use according to the label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### CONTAINER DISPOSAL:

**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill, incinerate or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

For First Aid, Precautionary Statements, and Directions for Use, see inside of the booklet.