

RESTRICTED USE PESTICIDE

Because pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and for only those uses covered by the Certified Applicator's certification.

FIFRA 24(c) Special Local Need Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Kerb® SC

EPA Reg. No. 62719-578

24(c) Special Local Need Registration SLN MI-120006

Weed Control in Direct Seeded Head Lettuce Grown in Muck Soils

(For Distribution and Use Only in the State of Michigan)

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2026

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read this SLN label and the label affixed to the container for Kerb® SC herbicide before applying.
- Use of Kerb SC according to this SLN label is subject to all use precautions and limitations imposed by the label affixed to the product container.

Direct Seeded Head Lettuce Grown in Muck Soils

General Information

Kerb SC is a selective herbicide for the preemergence control of certain grasses and annual broadleaf weeds in direct seeded head lettuce.

Kerb SC is a soil active herbicide that is absorbed through weed roots. In order to achieve good weed control, sufficient soil moisture from rain or irrigation is required following the treatment to move Kerb SC into the soil to the weed root zone.

Dosage and Weeds Controlled

For the control of susceptible weed species, apply Kerb SC at the rate of 9.6 to 14.4 pints (4 to 6 lb active ingredient) per acre. Kerb SC at the rate 9.6 pints per acre is effective for the control of common purslane in muck soils.

Kerb SC at the rate of 14.4 pints per acre will generally provide preemergence control of the following weeds when applied according to label directions:

common purslane (*Portulaca oleracea*)
common lambsquarters (*Chenopodium album*)

Kerb SC at the rate of 14.4 pints per acre will provide suppression to the following weeds and grass:
carpetweed (*Molluga verticillata*)
hairy nightshade (*Solanum sarrachoides*)
redroot pigweed (*Amaranthus retroflexus*)
barnyardgrass (*Echinochloa crusgalli*).

Dosage recommendations listed on this label are for surface broadcast applications. For banded treatments, the amount of Kerb SC used per acre should be reduced according to the following formula:

$$\frac{\text{Band width (in inches)}}{\text{Row width (in inches)}} \times \text{Rate Per Acre Broadcast} = \text{Amount Needed Per Acre Band Application}$$

Application Mixing and Equipment

Mix the specified amount of Kerb SC herbicide in clean water and apply uniformly in 30 to 50 gallons of water per acre. Use conventional ground spray equipment. Accurately calibrate spray equipment before each use.

Application Timing and Irrigation

For best control of susceptible weeds, applications of Kerb SC should be made immediately after planting operation, preemergence to the head lettuce and the weeds. The application should be followed immediately with an initial irrigation of 1 to 1-1/2 inches by overhead sprinkler equipment in order to move Kerb SC into the root zone of the germinating weeds.

Specific Use Restrictions

- Do not apply more than one application to each crop of head lettuce.
- Do not apply to head lettuce varieties which will be harvested less than 55 days after treatment.
- Do not plant back crops other than head lettuce for a period of 12 months following application.
- For use only for head lettuce grown on muck soils (muck soils should have an organic matter content of 20% or greater).
- Do not apply to leaf lettuce

™®Trademarks of Corteva Agriscience and its affiliated companies

R361-039

Accepted: 04/07/22

Replaces: R361-026