

Supplemental Label



Zidua® SC
Herbicide

For weed control in peanut

This supplemental label expires September 30, 2020, and must not be used or distributed after this date.

Active Ingredient:

pyroxasulfone: 3-[[[5-(difluoromethoxy)-1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl]methyl]sulfonyl]-4,5-dihydro-5,5-dimethylisoxazole 41.46%

Other Ingredients: 58.54%

Total: 100.00%

Contains 4.17 pounds of pyroxasulfone per gallon formulated as a water-based suspension concentrate

EPA Reg. No. 7969-374
CAUTION/PRECAUCION

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Zidua® SC herbicide** container label, EPA Reg. No. 7969-374, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Zidua SC** before applying.
- Use of **Zidua SC** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Zidua SC**.

Product Information

Zidua SC is a selective rate-dependent residual preemergence herbicide for controlling annual grasses, sedges, and annual broadleaf weeds. Refer to the **Zidua SC** container label for a complete list of weeds controlled.

Periods of dry weather following application of **Zidua SC** may reduce herbicidal effectiveness. **Zidua SC** must be activated by at least 1/2 inch of rainfall or irrigation before weed germination and emergence. When **Zidua SC** is not activated and weeds emerge, a labeled postemergence herbicide or shallow cultivation may be needed to control weed escapes. **Zidua SC** does not control emerged weeds.

Application Information

Zidua SC may be applied early postemergence to peanut for residual preemergence weed control.

Before applying to peanut, verify with your local seed company (supplier) the selectivity of **Zidua SC** on your variety to avoid potential injury.

Application Rate

Apply **Zidua® SC herbicide** alone, in tank mix, or sequentially in peanut at the residual rates are provided in **Table 1**.

Table 1. Residual Rates of Zidua SC in Peanut

Application Timing	Use Rate by Soil Texture ¹ (fl ozs/A)		
	Coarse	Medium	Fine
Early Postemergence	2.5 to 3.5	2.5 to 3.5	2.5 to 3.5

¹ Refer to container label for soil-texture groups.

Application Timing

Zidua SC may be applied in a single application or in sequential applications.

Early Postemergence Application

Apply **Zidua SC** at use rates specified in **Table 1** as a broadcast spray to peanut from “at-cracking” stage to first true leaf stage through beginning of pod development stage. **Zidua SC** provides residual control of weeds germinating after application. Weeds that are already emerged at the time of application must be controlled with cultivation, or tank mix or sequential application of another herbicide labeled for postemergence control of the target weeds in peanut. **Zidua SC** applications to emerged peanut may result in temporary leaf burn and stunting, but a reduction in peanut yield is unexpected. Tank mixes of **Zidua SC** with other crop protection products or adjuvants may significantly enhance this effect. Depending upon growing conditions, recovery from this injury begins immediately but may take several weeks for the injury to dissipate entirely.

Adjuvants may be applied with **Zidua SC** when making early postemergence applications.

Sequential Applications

If a sequential application program of **Zidua SC** is used (e.g. consecutive postemergence applications), the maximum combined rate of **Zidua SC** that may be applied in a cropping season (per year) is 8.25 fl ozs/A on all soils. Separate sequential applications by at least 14 days.

State-specific use in Texas in areas west of Interstate 35. Apply **Zidua SC** early postemergence at 2.5 fl ozs/A. Use of **Zidua SC** may result in growth suppression if heavy rainfall or irrigation (> 2 inches) occur after application. If a sequential application program of **Zidua SC** is used (e.g. consecutive postemergence applications), the maximum combined rate of **Zidua SC** that may be applied in a cropping

season (per year) is 5.0 fl ozs/A on all soils. Separate sequential applications by at least 21 days.

Crop-specific Restrictions

- **DO NOT** apply more than 3.5 fl ozs/A of **Zidua SC** (0.112 lb ai/A of pyroxasulfone) in a single application.
- **DO NOT** apply more than a maximum cumulative amount of 8.25 fl ozs/A of **Zidua SC** (0.269 lb ai/A of pyroxasulfone) per cropping season (per year).

Crop-specific Precautions

- **Zidua SC** applied early postemergence (at-cracking through first leaf stage) may result in temporary growth suppression in peanut if extreme conditions of high rainfall and extended periods of water-saturated soil occur during peanut early seedling development.
- There is no required (preharvest) interval between an early postemergence application of **Zidua SC** and harvest of peanut.

Tank Mixes

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Zidua SC may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products for a broader spectrum of control and/or control of emerged weeds. Refer to the tank mix product labels to confirm that the respective tank mix products are registered for use on peanut. Follow the adjuvant directions for the tank mix partner of **Zidua SC**.

- **Basagran® 5L herbicide**
- **Cadre® herbicide**
- **Outlook® herbicide**
- **Poast® herbicide**
- **Prowl® H2O herbicide**
- **Pursuit® herbicide**
- paraquat

Zidua SC may be tank mixed or applied sequentially with fungicide or insecticide products.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

***Basagran, Cadre, Outlook, Poast, Prowl, Pursuit, and Zidua** are registered trademarks of BASF.*

The  symbol is a registered trademark of Kumiai Chemical Industry Co., Ltd.

© 2017 BASF Corporation
All rights reserved.

007969-00374.20171130.**NVA 2017-04-489-0237**
Based on: NVA 2017-04-489-0046

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry