



2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA

DUPONT™ PREVATHON® INSECT CONTROL

EPA Reg. No. 352-844

FOR SUPPRESSION OF LESSER CORNSTALK BORER (*Elasmopalpus lignosellus*) IN PEANUT IN THE STATE OF TEXAS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

DuPont™ PREVATHON® insect control is recommended for suppression of lesser cornstalk borer (*Elasmopalpus lignosellus*) in peanut.

Apply PREVATHON® at 14 - 20 fluid ounces per acre (0.047 – 0.067 lb ai/acre) as a foliar spray for lesser cornstalk borer infestations later in the peanut growing season. Foliar applications should be made in sufficient water volume to achieve thorough coverage of peanut foliage and the soil area around the peanut main stem and growing limbs.

Applications of PREVATHON® by conventional foliar application methods should be followed with an overhead irrigation of 0.5 – 1.0 inches of water within 24 – 48 hours after application to enhance contact with lesser cornstalk borer larvae.

DuPont™ PREVATHON® may also be applied through overhead sprinkler irrigation systems, including the following; center pivot, end tow, hand move, lateral move, side roll, solid set and wheel line. The irrigation system used must provide uniform water distribution. Apply in no more than 0.2 inches of water per acre. For overhead sprinkler systems that are stationary, add the solution containing PREVATHON® to the irrigation water line and apply no more than 0.2 inches of water per acre. See the PREVATHON® Federal Section 3 label for complete guidance in making applications by overhead chemigation.

Peanut fields with a history of lesser cornstalk borer populations should be scouted weekly and PREVATHON® application made early when adult moths are active and laying eggs and when larvae are just beginning to hatch.

Apply when populations reach local established treatment thresholds to prevent crop damage.

Do not make more than two sequential applications of PREVATHON® before rotating to another registered insecticide having a different mode-of-action. See the RESISTANCE MANAGEMENT section of the PREVATHON® Section 3 Federal label for additional guidance on insecticide resistance management.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with such uses not specified by the label or DuPont 2(ee) instructions

© 2017 E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A.
All rights reserved.

PRECAUTIONS

Do not apply more than 60 fluid ounces of PREVATHON® or 0.2 lb ai chlorantraniliprole containing products per acre per year.

Do not make more than 4 applications per acre per crop. Minimum interval between treatments is 3 days.

Preharvest interval is 1 day.

The field re-entry interval is 4 hours.

Read and follow all applicable directions, restrictions, and precautions.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the DuPont product(s) covered by this Bulletin. DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by DuPont on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2017 E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A.
All rights reserved.