



FIFRA Section 24 (c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

ACEPHATE 97UP INSECTICIDE

EPA Reg. No. 70506-8

EPA SLN No. ID-170005

ACTIVE INGREDIENT	BY WT.
Acephate: (O,S-Dimethyl acetylphosphoramidothioate).....	97%*
OTHER INGREDIENTS	3%
TOTAL	100 %

*U.S. Patent Nos. 6,752,943 and 6,875,381

CAUTION

FOR INSECT CONTROL IN ALFALFA GROWN FOR SEED ONLY

This label is valid until December 31, 2020 or until otherwise amended, withdrawn, cancelled or suspended.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This FIFRA Section 24 (c) Special Local Need (SLN) label and the federal label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label. Read the entire Directions for Use before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical resistant gloves made of any waterproof material and shoes plus socks.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds.

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash-waters.

Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Eliminations System (NPDES) permitted and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

USE RESTRICTIONS

Do not apply this product through any type of irrigation system.

No portion of this alfalfa seed crop may be used or distributed for human food or animal feed purposes.

Alfalfa Grown For Seed For Control of Lygus Bug

PESTS CONTROLLED	RATE	APPLICATION INSTRUCTIONS	COMMENTS
Lygus Bug	Use 1 lb (0.97 lb ai) of Acephate 97UP per acre per application. Do not exceed 2 lbs (1.94 lbs ai) of product per acre per year.	Ground – apply in 10 to 60 gallons of water. Air – apply in 5 to 10 gallons of water per acre. Make one repeat application at no less than 7 day spray interval, if necessary to maintain control.	Applications may be made up to 3 days prior to placing bees in or around the field. Do not make more than 2 applications per acre per year. The REI is 24 hours. Some data indicate that Leafcutter bees and Alkali bees may be more sensitive to acephate exposure than other tested species.