



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA® ON GARLIC

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

REGALIA®

EPA Reg. No. 84059-3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROPS	DISEASES	APPLICATION METHOD	RATES	USE RECOMMENDATIONS
Bulb Vegetables Garlic	Botrytis Leaf Blight (<i>Botrytis squamosa</i>) Botrytis Neck Rot (<i>Botrytis</i> spp.) Downy Mildew (<i>Peronospora</i> spp.) Onion Purple Blotch (<i>Alternaria porri</i>) Powdery Mildew (<i>Erysiphe</i> spp.)	Foliar (ground)	1-4 quarts per acre	For foliar applications, apply this product preventatively in 50 -100 gallons of water per acre. Do not exceed 1% v/v of the applied solution. Repeat applications at 7-14 day intervals. Under moderate to heavy disease pressure tank-mix this product with another fungicide.
	<i>Pythium</i> spp. <i>Rhizoctonia</i> spp.	Soil drench	1-2 quarts per 100 gallons	For soil drench applications, apply this product at a concentration of 1-2 quarts per 100 gallons of water, and at a rate to thoroughly soak the growing media and root zone. Make an initial application of this product during or shortly after transplant to reduce transplant shock, suppress soil-borne diseases and improve root growth. Multiple drench applications can be made on a 10-14 day interval.

Bulb Vegetables Garlic	<i>Pythium</i> spp. <i>Rhizoctonia</i> spp.	In-furrow	1-2 quarts per acre 1.2 – 3.7 fluid ounces per 1000 foot of row	For in-furrow applications, at planting apply this product as an in-furrow spray at the rate of 1-2 quarts per acre or 1.2 – 3.7 fluid ounces per 1000 foot or row. Apply this product in 5-15 gallons of water per acre so as the spray is directed into the seed furrow just before the seeds are covered.
	<i>Pythium</i> spp. <i>Rhizoctonia</i> spp	Chemigation	1-4 quarts per acre	For chemigation applications, apply this product at 1-4 quarts per acre immediately after transplant and at 14 day intervals or begin 14 days after transplant when plant dip or in-furrow applications are used.
	<i>Pythium</i> spp. <i>Rhizoctonia</i> spp	Plant dip	1-2 quarts per 100 gallons	For plant dip applications for improved plant health and suppression of soil-borne diseases, apply this product in a 0.25-0.5% v/v suspension (1-2 quarts per 100 gallons water) as a pre-plant dip immediately prior to transplanting.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

This 2(ee) recommendation expires December 31, 2019.

Regalia[®] is a registered trademark of Marrone Bio Innovations, Inc.
 © Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.
 1540 Drew Ave.
 Davis, California 95618
 Telephone (530) 750-2800
www.marronebioinnovations.com