

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Stinger\*

EPA Reg. No. 62719-73

24(c) Special Local Need Registration EPA SLN No. MT-99-0011  
For Distribution and Use Only in the State of Montana

**For Control of Canada Thistle and Perennial Sowthistle  
Infesting Canola Grown Only for Seed**

**NOTICE:** A nonfood use agreement must be signed with the Montana Department of Agriculture prior to purchase of this product for use in seed canola.

**Do not feed or graze canola fields following application of Stinger\* herbicide. Do not cut canola for forage. Do not use harvested seed for oil production. Processed seed must be labeled "Not for human or animal consumption." All canola seed screenings must be disposed of in such a way that they cannot be distributed or used for food or feed.**

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Stinger before applying. Carefully follow all precautionary statements and applicable use directions.
- In additions to restrictions and limitations imposed by this supplemental labeling, use of Stinger is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger.

### Directions for Use

Stinger may be applied to control Canada thistle or perennial sowthistle infesting canola.

**Application Method:** Apply as a broadcast spray in a total spray volume of 10 or more gallons of water per acre using ground equipment.

#### Broadcast Rates and Application Timing:

Weeds Controlled	Application Rate (pt/acre)	Application Timing
Canada thistle	Light infestation - 1/3 Moderate to heavy infestation - 1/2	Apply when the majority of thistle plants have emerged, and thistles are at least 6 to 8 inches in diameter or height up to bud stage. Extreme growing conditions (such as drought or near freezing temperatures) prior to, at, and following the time of application may reduce weed control.
Perennial sowthistle	1/2	

#### Restrictions:

- Maximum number of applications: 1
- Carefully follow rotational crop restrictions and other use precautions and limitations on the product label for Stinger.
- **Preharvest Interval:** Do not apply within 72 days of harvest.

\*Trademark of Dow AgroSciences LLC

D06-043-013

Accepted: 04-27-99

Initial printing.