

Andersons Golf Products®

24-0-18 with Fungicide VII

• Provides systemic prevention and control of turfgrass diseases including benzimidazole tolerant dollar spot

• Provides controlled release nitrogen feeding

ACTIVE INGREDIENT:	
Triadimefon†	0.62%
OTHER INGREDIENTS:	99.38%
†CAS No. 43121-43-3	Total 100.00%

EPA Reg No. 9198-190
EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code on bag.

Guaranteed Analysis	
Total Nitrogen (N)	24.00%
9.72% Urea Nitrogen	
5.88% Water Insoluble Nitrogen	
8.40% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	18.00%
Sulfur (S)	6.12%
6.12% Combined Sulfur (S)	

Plant nutrients derived from urea, methylene ureas and potassium sulfate.
*8.40% Slowly available nitrogen from methylene ureas.

F292

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-ap.com>

U.S. Patent Nos. 5,102,440 and 4,378,238

Attention: This product contains a chemical known to the State of California to cause birth defects or other reproductive harm.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)
Loaders, applicators, flaggers and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks, and
- Chemical-resistant gloves made of any waterproof material

User Safety Requirements
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations
Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
If swallowed:	Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If inhaled:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.	

Environmental Hazards: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift from areas treated. Drift and runoff from treated areas may be hazardous to organisms in neighboring areas. Apply this product only as specified on this label.

This product may contaminate water through runoff. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

DIRECTIONS FOR USE
It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

For use on turfgrass areas consisting of Kentucky bluegrass, annual bluegrass, bentgrass, perennial ryegrass, fine fescue, bermudagrass or mixtures where these grasses predominate.

Precautions
Do not enter or allow others to enter the treated area (except those involved in watering-in) until watering-in is complete and the surface is dry.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Application Restrictions for Turfgrass Uses
Use on turfgrass is limited to golf course turf only. Use on turfgrass is prohibited on all other residential and commercial sites.

Maximum application rate is 2.7 lbs. ai/A
Maximum annual application rate is 5.4 lbs ai/A
Minimum retreatment interval is 14 days

Application to golf courses, including tees, greens, fairways and roughs is permitted only if the turfgrass is 2.5 inches or less in height.

TO PREVENT DOLLAR SPOT, ANTHRACNOSE AND RUST (*Puccinia spp.*), apply to moist or dry foliage at the DIRECTED RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE DOLLAR SPOT, ANTHRACNOSE AND RUST (*Puccinia spp.*), apply at the DOUBLE RATE to moist or dry foliage at 2 week intervals. When the disease activity is under control, apply at the DIRECTED RATE at two week intervals.

TO PREVENT BROWN PATCH, apply to moist or dry foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE BROWN PATCH, use a non-fertilizer fungicide such as Andersons Golf Products Fungicide VII to limit the amount of fertilizer applied since high levels of nitrogen applied to turf during or just prior to the onset of hot, humid weather may increase the severity of brown patch. Follow with the DIRECTED RATE of either product once the disease is controlled.

NOTE CAREFULLY
To avoid product pickup, lightly irrigate treated areas soon after application.

On short cut bentgrass (less than 1/2") when temperatures are above 80°F, apply only to dry foliage.

Storage and Disposal Do not contaminate water, food, or feed by storage or disposal. STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing. Protect from water and extreme heat. PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.
--

Spreader Settings				
These spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product. 27.72 lbs feeds 22,000 sq. ft. (0.50 acre) with 0.30 lbs. N/1,000 sq. ft. at the DIRECTED RATE 27.72 lbs feeds 11,000 sq. ft. (0.25 acre) with 0.60 lbs. N/1,000 sq. ft. at the DOUBLE RATE				
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			DIRECTED RATE	DOUBLE RATE
Andersons Model 2000 / AccuPro 2000, (cone 8) Andersons 2000 SR / SR-2000 (cone 8)	3 mph	10 feet	H	I 1/2
Andersons SSD / SS-2 Drop	3 mph	overlap wheels	2 1/4	3
Scotts R-8, R-8A (cone 8)	3 mph	10 feet	G	H 1/2
Lesco 020093 (pattern 0.0)	3 mph	8 feet	B 1/2	C 1/2
Lesco 705698 (pattern 0.0)	3 mph	8 feet	7	9

Note: These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

IMPORTANT: READ BEFORE USE:
Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:
Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:
THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Product of U.S.A.

Manufactured by:
The Andersons Lawn Fertilizer Division, Inc.
P.O. Box 119, Maumee, OH 43537
www.andersonsturf.com

Andersons Golf Products is a registered trademark of The Andersons, Inc.

SGN 100
Net Wt. 27.72 Lbs. (12.57 kg)



AGC8578.1
G12 (AGCBLKP1626.1B)