

SUPPLEMENTAL LABELING

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**
TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS
WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR, EXCEPT THAT UNCERTIFIED
PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina, 27419-8300
SCP 1652A-S1A 0822

PARAQUAT DICHLORIDE	GROUP	22	HERBICIDE
---------------------	-------	-----------	-----------

Gramoxone® SL 3.0
HERBICIDE

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

Active Ingredients:

Paraquat dichloride*	43.9%
Other Ingredients:	56.1%
Total:	100.0%

*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid containing 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.

Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



DANGER / POISON
PELIGRO / VENENO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)
See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



**DIRECTIONS FOR USE
RESTRICTED USE PESTICIDE**

Citrus Hybrids, CA Only

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Clementines Orange, Temple		Tangelos Tangors	
Target Weed	Rate pt/A	Application Timing	Use Directions
Refer to Section 7.0.	1.7 – 2.7	Postemergence (Directed Spray)	<p>For use in California Only.</p> <p>Apply as a ground directed spray in a minimum of 10 gal/A of spray volume.</p> <p>The recommended spray volume is minimum, and spray volumes should be increased as necessary to obtain complete coverage of the target weed or plant without runoff from the foliage.</p> <p>Make ground applications when the wind velocity favors on target product deposition, (approximately 3-10 mph). Do not apply when wind velocity exceeds 10 mph.</p> <p>See Section 4.4.4 of the Gramoxone SL 3.0 for adjuvant recommendations.</p>
Tank Mix Options:			
<ul style="list-style-type: none"> • Gramoxone SL 3.0 can be tank mixed with other herbicides labeled for use on citrus hybrids applied by the same methods and at the same timings. Refer to Section 4.4.2 for tank-mix precautions in the Gramoxone SL 3.0 label. 			
Resistance Management:			
<ul style="list-style-type: none"> • Refer to Section 3.1. 			
Precaution:			
<ul style="list-style-type: none"> • Use a shield or wrap plant when spraying around young trees. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1. Refer to Section 6.1 for additional product use restrictions. 2. Maximum Single Application Rate: 2.7 pt/A 3. REI: 48 hours 4. Minimum Application Interval: 7 days 5. Maximum Annual Rate: 13.3 pt/A <ol style="list-style-type: none"> a. DO NOT exceed 5.0 lb cation/A/year of paraquat-containing products. 6. DO NOT make more than 5 applications per year. 7. DO NOT allow spray to contact green stems (except suckers), fruit or foliage. 8. DO NOT apply when weather conditions favor drift from treated areas. 9. DO NOT apply by ground equipment within 25 ft or by air within 75 ft of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds. 10. DO NOT apply this product through any type of irrigation system. 11. DO NOT graze treated areas. 12. DO NOT feed cover crops grown in treated areas to livestock. 13. Pre-Harvest Interval (PHI): Harvest at normal crop maturity. 			

Gramoxone®, and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2022 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina, 27419-8300