



The Go To Company

P.O. Box 5569 • Yuma, AZ 85366 • Toll Free: 800.883.1844 ext. 2 • www.gowanco.com

**FIFRA §24(c) SPECIAL LOCAL NEED REGISTRATION**  
**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON**  
**FOR CONTROL OF SPOTTED WING DROSOPHILA IN CANEBERRIES**

**GOWAN MALATHION 8 FLOWABLE**  
**AGRICULTURAL INSECTICIDE**

EPA Reg. No. 10163-21

SLN No. OR-130011

**This label is valid until December 31, 2022**  
**or until otherwise amended, withdrawn, canceled, or suspended**

ACTIVE INGREDIENT:	% By Wt.
Malathion (O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate):	79.5%
INERT INGREDIENTS	20.5%
	TOTAL 100.0%

Contains Petroleum Distillates  
Contains 8 lbs. Malathion per gallon

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions and precautions on the EPA registered label and this label for Gowan Malathion 8 Flowable (EPA Reg. No. 10163-21).
- This labeling must be in the possession of the user at the time of pesticide application.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to aquatic organisms, including fish and invertebrates. This product may contaminate water through drift of spray in wind. This product has a high potential for runoff after application. Use care when applying in or to an area which is adjacent to any body of water, and do not apply when weather conditions favor drift from target area. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

**DIRECTIONS FOR USE**

CROP	REI	RATE (PTS/ACRE)	PEST	RESTRICTIONS
BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES	12 hours	Up to 2 (2.0 lbs ai/acre)	Spotted Wing Drosophila	The maximum application rate is 2.0 pints of product per acre; the maximum number of applications per year is 4; and the minimum retreatment interval is 7 days.  Do not exceed a total maximum use rate of malathion from all sources of 8 lbs. ai per acre per year.  Do not apply within 1 (one) day of harvest.

**IMPORTANT:** This product is sold subject to the Conditions of Sale and Warranty and Liability Limitations set forth on the container label.

**24(c) REGISTRANT:** Gowan Company  
P.O. Box 5569  
Yuma, AZ 85366-5569