

## FIFRA 2(ee) Recommendation

EPA Reg. No.: 279-3618

## FOR DISTRIBUTION AND USE ONLY IN TEXAS

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation is valid until December 31, 2028 or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED SUBJECT TO THE TERMS AND CONDITIONS OF THE PRODUCT LABELING.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

## Addition of nematode pests suppressed by Zironar® Biofungicide/Bionematicide

CROP	EARLY SEASON PROTECTION OF SEEDLINGS AGAINST INJURY FROM	APPLICATION RATE
Corn Field Corn, Popcorn, Sweet Corn, and Corn grown for seed	Dagger Nematode (Suppression) Pin Nematode (Suppression) Spiral Nematode (Suppression) Stunt Nematode (Suppression) Stubby-root Nematode (Suppression) Lance Nematode (Suppression) Needle Nematode (Suppression) Sting Nematode (Suppression)	6 to 12 fl oz/A Apply in a minimum of 3 gallons of water per acre.
Cotton  Directions For Use	Lance Nematode (Suppression) Sting Nematode (Suppression)	6 to 12 fl oz/A Apply in a minimum of 3 gallons of water per acre.
Apply in the open furrow in a narrow band at the seeding depth or as a T-band over the open furrow during planting.		

R-4925 121624 10-28-22

