



Raxil[®]

PRO MD

Net
Contents:

2.5 Gallons



Bayer SeedGrowth™

ACTIVE INGREDIENTS:

Prothioconazole: 2-[2-(1-Chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-1,2-dihydro-3H-1,2,4-triazole-3-thione	1.47%
Tebuconazole: (Alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol)	0.29%
Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester	0.59%
OTHER INGREDIENTS:	97.65%
TOTAL:	100.00%

Contains 0.128 pounds Prothioconazole, 0.025 pounds Tebuconazole and 0.052 pounds Metalaxyl per U.S. gallon.

EPA Reg. No. 264-1072

**KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO**

See Additional Precautionary Statements

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

Produced for:
Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
RAXIL is a registered trademark of Bayer.
©2019 Bayer CropScience

160127CV2 02/19

US61380807C

FIRST AID

IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything to an unconscious person.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
<p>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577. Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	
<p>NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Causes irreversible eye damage. Harmful if swallowed, inhaled, or absorbed through skin. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Avoid contact with skin and breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Workers involved with treating the seed (e.g. connecting and disconnecting hoses and transfer pumps, mixing, equipment calibration, etc.) and other exposed to the concentrate, and cleaners/repairers of seed treatment equipment must wear a long-sleeve shirt and long pants, shoes plus socks, protective eyewear (goggles, face shield, or safety glasses) and chemical resistant gloves (such as natural rubber, follow the instructions for category A on an EPA chemical-resistance category selection chart).

Baggers and bag sewers must wear a long-sleeve shirt and long pants, and shoes plus socks.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothes.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish, aquatic invertebrates, and freshwater/estuaries/marine aquatic plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

Cover or incorporate spilled treated seed.

Groundwater Advisory: Prothioconazole-desthio (a degradate of prothioconazole), metalaxyl and tebuconazole are known to leach through soil into ground water under certain conditions as a result of label use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

PRODUCT INFORMATION

Raxil® PRO MD is a combination of the systemic fungicides, prothioconazole, tebuconazole and metalaxyl which aids in the control or suppression of the following seed, seedling and soilborne diseases of wheat, barley, oats, and triticale: stinking smut, flag smut, loose smut, covered smut (common bunt), early season *Septoria* disease complex, general seed rots, *Pythium* damping-off, early season *Rhizoctonia* root rot, early season common root rot, seedborne *Fusarium* scab, barley stripe, early season *Fusarium* foot rot, early season suppression of powdery mildew (7.5 fluid ounce rate only) and rust.

In California – Registered only for use in commercial seed treating establishments. Not allowed for “on-farm” use.

APPLICATION DIRECTIONS

Apply this product only as specified on this label. Use 5.0 – 7.5 fluid ounces of Raxil PRO MD per 100 pounds of seed. Applications must be made using standard slurry or mist-type seed treatment equipment. In some instances, a chemical slurry dilution may enhance coverage. A 1 part Raxil PRO MD with 1 part water dilution may be used. This product is for commercial or on-farm application. Do not apply this product with a hopper box, planter box, slurry box, or any other at-plant methods. Uniform application to seed is necessary to ensure seed safety and best disease control. Only use seed that is sound and well cured prior to treatment. Consult your area agronomist regarding planting depth recommendations for the particular variety to be planted.

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with Raxil PRO MD must be labeled with the following statements:

This seed has been treated with Raxil PRO MD, which contains 1.47% of Prothioconazole, 0.29% of Tebuconazole, and 0.59% of Metalaxyl.

- Do not use treated seed for food, feed, or oil purposes. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Barley, wheat, oats, and triticale green forage may be grazed or harvested for hay 31 days after seeding.

(continued)

- Store treated seeds away from food and feedstuffs.
- Do not allow children, pets, or livestock to have access to treated seed.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed. Leftover treated seed may be buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Treated seed must be adequately covered with soil at planting.
- Dispose of seed packaging or containers in accordance with local requirements. Do not use empty seed bags for any other purpose.
- To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's recommendations. Surplus seed or empty seed containers should be stored or disposed according to local federal regulations.
- Treated at manufacturer's recommended rate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment.

PESTICIDE STORAGE

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat.

PESTICIDE DISPOSAL

Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING

Dilutable Seed Treatment Products in Non-Refillable Plastic Containers

Nonrefillable container. Do not reuse or refill this container. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781), or rinse and either recycle or dispose of the container as follows:

Liquid dilutables in containers small enough to shake (5 gallons or less)

Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer

NET CONTENTS: 2 1/2 GALLONS

Raxil® PRO MD

ACTIVE INGREDIENTS:

Prothioconazole: 2-[2-(1-Chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-1,2-dihydro-3H-1,2,4-triazole-3-thione	1.47%
Tebuconazole: (Alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol)	.029%
Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester	.59%
OTHER INGREDIENTS:	.9765%
TOTAL:	100.00%

Contains 0.128 pounds Prothioconazole, 0.025 pounds Tebuconazole and 0.052 pounds Metalaxyl per U.S. gallon.

EPA Reg. No. 264-1072

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

See Additional Precautionary Statements

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Causes irreversible eye damage. Harmful if swallowed, inhaled, or absorbed through skin. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Avoid contact with skin and breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment.

PESTICIDE STORAGE

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat.

PESTICIDE DISPOSAL

Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING

Dilutable Seed Treatment Products in Non-Refillable Plastic Containers

Nonrefillable container. Do not reuse or refill this container. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781), or rinse and either recycle or dispose of the container as follows:

Liquid dilutables in containers small enough to shake (5 gallons or less)

Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167

RAXIL is a registered trademark of Bayer.
©2019 Bayer CropScience

US61380807C

160127Cv2 02/19



(01) 0 0785740 34126 1