

**RESTRICTED USE PESTICIDE  
DUE TO ACUTE TOXICITY**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**syngenta.**

**INTIRA**

**Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS**

**Gramoxone SL 2.0**

**For Preplant / Preemergence Control of Weeds in Peanuts**

**EPA Reg. No. 100-1431  
EPA SLN No. TX-190011**

**This label expires and must not be distributed or used in accordance with this  
SLN registration after December 31, 2024**

**KEEP OUT OF REACH OF CHILDREN**



**DANGER / POISON  
PELIGRO / VENENO**

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- Failure to follow the Directions for Use and Precautions on the EPA-registered Gramoxone SL 2.0 label may result in poor pest control, crop injury, or illegal residues.

<b>PEANUTS</b>		
<b>Rate pt/A</b>	<b>Application Timing</b>	<b>Use Directions</b>
1.9 – 3.75	Preplant / Preemergence	<p>Apply to burndown weeds in peanut fields preplant / preemergence.</p> <p>Apply as a ground broadcast application in a minimum of 10 gal/A of spray volume. Increase carrier volume for dense weed populations and larger weeds</p> <p>Use the higher rate on dense populations and/or on larger or harder to control weeds. Apply when weeds are actively growing and between 1" to 6" in height. Vegetation 6" or taller may not be controlled.</p> <p>Always add either: 1) Nonionic surfactant containing 80% or more surface agent at 0.25% v/v (2 pints per 100 gallons), of the finished spray volume, <b>OR</b> 2) crop oil concentrate (non phytotoxic) containing 15 to 20% approved emulsifier, at 1% v/v (1 gallon per 100 gallons) of the finished spray volume.</p> <p><b>Tank Mix Options:</b> Gramoxone SL 2.0 may be tank-mixed with other herbicides registered for this use for improved weed control.</p>
<p><b>Use Restrictions:</b></p> <ol style="list-style-type: none"> <li><b>Maximum Single Application Rate:</b> 3.75 pt/A.</li> <li><b>Maximum Annual Rate:</b> 7.5 pt/A.  <b>DO NOT</b> exceed 1.9 lb ai/A/year of paraquat-containing products for preplant/preemergence uses.  <b>DO NOT</b> exceed 1.9 lb ai/A/year of paraquat-containing products for all combined uses.</li> <li><b>DO NOT</b> make more than 2 preemergence applications per year.</li> <li><b>DO NOT</b> apply this product by air.</li> <li><b>DO NOT</b> allow livestock to graze in treated areas.</li> <li><b>Pre-Harvest Interval (PHI):</b> NA.</li> </ol>		

©2019 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company.

24(c) Registrant:  
 Syngenta Crop Protection, LLC  
 P. O. Box 18300  
 Greensboro, NC 27419-8300

Label Code: TX1431178AA0519