

Product Bulletin

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Opensight®

EPA Reg. No. 62719-597

2(ee) Recommendation†

For Distribution and Use Only in the State of Texas

This product bulletin expires December 31, 2023, and must not be used or distributed after this date.

Control of Susceptible Weeds Pre or Post Planting of Certain Annual and Perennial Grasses On Rangeland, Permanent Grass Pastures, Conservation Reserve Program (CRP) Acres, Natural Areas, and Grazed Areas in and Around These Sites

ATTENTION

† This recommendation is made as permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions, and limitations on the product label attached to the container for this product.

Directions for Use

Opensight provides preemergence and postemergence control of susceptible broadleaf plants on rangeland, permanent grass pastures, CRP acres, natural areas, and grazed areas in and around these sites. See the main label for Opensight for a list of susceptible weeds.

Precautions

Post emergence applications on newly emerged tall fescue, or newly emerged or established annual ryegrass, timothy, sorghum-sudangrass, teff, crabgrass, and pearl millet are not recommended. Unacceptable injury or death of these grasses may occur with Opensight applications.

Pre-plant applications for control of susceptible emerged weeds

Apply Opensight at a maximum of 2.0 oz/acre with a non-ionic surfactant to control susceptible weeds no sooner than 30 days prior to planting tall fescue, orchardgrass, timothy, and annual ryegrass. Opensight can also be used at 2.0 oz/acre prior to planting sorghum-sudangrass, teff, crabgrass, and pearl millet, but a minimum 45-day interval must be observed between the application and planting.

Post planting applications for control of susceptible emerged weeds

Apply Opensight at a maximum of 2.0 oz/acre with a non-ionic surfactant to orchardgrass after seedling plants have at least began tillering (multiple stems from single crown).

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R372-028

Issued: 08/10/2020

Replaces R372-026