



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

ALIETTE® WDG Fungicide

For Aerial Application to Strawberries

EPA Reg. No. 264-516

EPA SLN. No. CA-970008

24(c) Supplemental Label

FOR USE AND DISTRIBUTION ONLY IN THE STATE OF CALIFORNIA

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

Crop/Site:	Strawberry / California
Target Pest:	Red Stele (<i>Phytophthora fragariae</i>); Leather Rot (<i>Phytophthora cactorum</i>)
Rate:	2.5 - 5.0 lbs product per acre
Dilution Rate:	Minimum of 10 gallons per acre
Method of Application:	Aerial
Frequency and Timing of Application:	Do not apply more than 30 lbs per season. For Red Stele begin applications 14-21 days after planting and continue on a 30-60 day interval as long as conditions favor disease development. For Leather Rot begin applications between 10% bloom and early fruit set and continue on a 7-14 day interval as long as conditions are favorable for disease development. The shorter time intervals and higher rates should be used when disease pressure is high.
Field Reentry Safety Interval:	Refer to the EPA registered label for ALIETTE® WDG Fungicide (EPA No. 264-516) for safety and re-entry information.

Other Requirements:

Note: Bayer CropScience has determined that phytotoxicity from solubilized copper may occur if products containing copper are tank-mixed with ALIETTE® WDG or if unbuffered ALIETTE® WDG is applied to foliage with copper residues. Do not tank-mix with copper fungicides.

When applied prior to or after copper fungicides, the pH of ALIETTE® WDG should be raised to 6.0 or above with the addition of an alkaline buffer such as Potassium Carbonate (3 lbs Potassium Carbonate to 5 lbs ALIETTE® WDG) or DiAmmonium Phosphate (5 lbs DiAmmonium Phosphate to 5 lbs ALIETTE® WDG).

Adjuvants which enhance pesticide penetration may cause phytotoxicity when mixed with ALIETTE® WDG.

Mixing ALIETTE® WDG with surfactants or foliar fertilizers is not recommended. Strawberry sensitivity to a particular combination should be determined by spraying a few plants. Evaluate 3-7 days later for adverse effects.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com