



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF WASHINGTON For Control of Pear Psylla Nymphs and Mites

GROUP	21A	ACARICIDE
-------	-----	-----------

Nexter[®] SC Miticide/Insecticide

EPA Reg. No 81880-28-10163 / EPA SLN NO. WA-170002b

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2021.

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

WSDA'S AQUATIC ADVISORY: This pesticide is toxic to fish and aquatic invertebrates. Nexter SC Miticide/Insecticide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

WSDA CONTAINER DISPOSAL GUIDANCE: Completely empty the contents of the pesticide container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

POLLINATOR PROTECTION: This product is highly toxic to bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Apply Nexter SC Miticide/Insecticide between late evening and early morning only.

CROP	RATE	PEST	COMMENTS
Pears (including oriental)	17 – 25 ounces / acre	Pear Psylla, European Red Mite, Twospotted Spider Mite, Pear Rust Mite	Apply Nexter SC using conventional dilute or concentrate ground sprayers calibrated to deliver enough water to ensure thorough coverage of foliage. Spray volume will vary with the size and number of trees per acre and foliage density. Nexter SC must be applied to each row for maximum coverage. Pear psylla numbers can increase rapidly; begin application before damaging numbers are reached. Applications may be made early from cluster bud through petal fall to control eggs, early pear psylla instars and mobile mites. Nexter SC is also effective when applied after petal fall as mite and psylla populations begin to build.
RESTRICTIONS / PRECAUTIONS: <ul style="list-style-type: none"> • Do not apply within 28 days of harvest (PHI). • Do not make more than 1 application per season. • Do not apply by air. • For use under this SLN label do not apply this product through any type of irrigation system. • Drift: Do not apply Nexter SC Miticide/Insecticide when weather conditions favor drift to surface water. Do not apply within 110 feet upwind of surface water or when windspeed is above 8 mph. Do not apply during a temperature inversion. 			

Nexter[®] is a registered trademark of Nissan Chemical Industries, Ltd.

24(c) Registrant: Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Updated 2-16-17