

**GROUP 4 HERBICIDE**

# FIFRA Section 24(c) Special Local Need (SLN) Label

**FOR DISTRIBUTION AND USE ONLY WITHIN OREGON**



## WEEDAR® 64 Broadleaf Herbicide

EPA Reg. No.: 71368-1

SLN No.: OR-010016

**SLN Label Expiration Date: December 31, 2020**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

### FOR CONTROL OF TALL WEEDS IN CRANBERRY BOGS

**ACTIVE INGREDIENT:**

2,4-Dichlorophenoxyacetic acid, dimethylamine salt\* ..... 46.8%

**OTHER INGREDIENTS:** ..... 53.2%

**TOTAL** ..... 100.0%

\*2,4-Dichlorophenoxyacetic acid equivalent 38.9% by weight or 3.8 pounds per gallon. Isomer specific by AOAC method No. 978.05

**KEEP OUT OF REACH OF CHILDREN**

**DANGER / PELIGRO**

PRECAUCION AL USUARIO: Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840**

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

**For Terrestrial Uses:** Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This product contains a chemical with properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**For Aquatic Uses:** Fish breathe dissolved oxygen in the water and decaying weeds also use oxygen. When treating continuous, dense weed masses, it may be appropriate to treat only part of the infestation at a time. For example, apply the product in lanes separated by untreated strips that can be treated after vegetation in treated lanes has disintegrated. During the growing season, weeds decompose in a 2 to 3 week period following treatment. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Waters having limited and less dense weed infestations may not require partial treatments.

Do not contaminate water used for irrigation or domestic purposes (except as specifically listed on this label) especially in areas where grapes, cotton, tomatoes or other susceptible plants are grown.

Do not treat irrigation ditches in areas where water will be used to overhead (sprinkler) irrigate susceptible crops especially grapes, tomatoes, tobacco, and cotton.

Do not apply this product directly to, or permit to drift onto cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants which are susceptible to 2,4-D herbicide. Do not apply near susceptible plants since very small quantities of the 2,4-D will cause severe injury during the growing or dormant periods. Crops contacted by this product sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

**MIXING AND LOADING:** Most cases of ground water contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent ground water contamination.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### GENERAL INFORMATION

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this product only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult Nufarm salesperson, State Agricultural Extension Service or qualified crop consultant for advice.

Be sure that use of this product conforms to all applicable laws, rules and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator should become familiar with these laws, rules or regulations and follow them exactly.

To be used in connection with WEEDAR<sup>®</sup> 64 Broadleaf Herbicide for control of the following weeds:

Asters	( <i>Aster</i> spp.)
Dewberries	( <i>Rubus</i> spp.)
Goldenrod	( <i>Solidago</i> spp.)
Hardhack	( <i>Spirea tomentose</i> L.)
Leatherleaf	( <i>Chamaedaphne calyculata</i> L.)
Loosestrife	( <i>Lysimachia terrestris</i> L.B.S.P.)
Ragweed	( <i>Ambrosia artemisiifolia</i> L.)
Red maple	( <i>Acer rubrum</i> L.)
Saw briar	( <i>Smilax glauca</i> )
St. Johnswort	( <i>Hypericum</i> spp.)
Sweetgum	( <i>Liquidambar styraciflua</i> )
Sweet pepper bush	( <i>Clethra alnifolia</i> L.)
Three square	( <i>Scirpus aminicanus</i> )

Best results when used in late June and July. **Preharvest interval is 30 days.**

Apply with a wooden frame or similar device, shaped like a hockey stick, with its lower member wrapped with several thicknesses of Turkish toweling (or other suitable material). Apply by soaking the toweling in one part WEEDAR® 64 herbicide to two parts water. Then with swabbed portion of the stick horizontal, wave left and right above the cranberry vines, wiping small quantities of the herbicide onto tall weeds above the crop level. **Do not allow pesticide to drip or touch cranberry vines.** Apply only once per year.

**PRECAUTIONS AND RESTRICTIONS**

Preharvest interval is 30 days. Apply only once per year.  
Do not exceed 2.5 pints per acre per application.

**NOTICE**

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

**Manufactured For Nufarm, Inc.  
11901 S. Austin Avenue, Alsip, IL 60803**

Weedar is a registered trademark of Nufarm, Inc.