



BISON™

Supplemental Label

CONSERVATION RESERVE PROGRAM (CRP) AREAS AND GRASSES GROWN FOR SEED PRODUCTION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT:

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Personal protective equipment and reentry intervals on the basic label must be followed.

Directions For Use:

CONSERVATION RESERVE PROGRAM AREAS (CRP) BISON RECOMMENDATIONS

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
BISON	1 - 2 pints/A	Apply to grasses from the 3-leaf stage	Refer to the GENERAL WEED LIST for a listing of susceptible broadleaf weeds. Apply 1 pint/ A to MOST SUSCEPTIBLE and 1 1/2 - 2 pints/ A to SUSCEPTIBLE broadleaf weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first..

RESTRICTIONS AND PRECAUTIONS: CRP AREAS

- Do not allow livestock to graze in treated areas or feed grass to livestock.
- If legumes are included in CRP area planting, severe injury to the legumes could result from treatment with BISON.

GRASSES GROWN FOR SEED OR SOD PRODUCTION
BISON RECOMMENDATIONS
Seedling and Established Grasses

PRODUCT	RATE/A	APPLICATION TIMING AND SPECIFIC COMMENTS		
		RATE/1000 sq. ft	CROP	WEEDS
BISON	1 - 2 pints/A	0.375 - 0.75 fl. oz.	<p>Apply to established and newly seeded grasses grown for seed or sod production before the boot stage.</p> <p>Established grasses tolerant to BISON include bentgrasses, Kentucky bluegrass, fescues, ryegrasses, Bermuda grass, St. Augustine grass and zoysiagrass. BISON may also be used on seedling grasses such as Merion, Park, Delta, or common Kentucky bluegrasses, Pennlawn, chewings, Illahee, or Alta fescues, orchard grass, Highland, Seaside, or Astoria bentgrasses, perennial ryegrasses, Bahiagrass and zoysiagrass.</p>	<p>Refer to the GENERAL WEED LIST for a listing of susceptible broadleaf weeds.</p> <p>Optimal control will be obtained when weeds are treated in the seedling stage (less than 4-leaf stage, 2 inches in height, or 1 inch in diameter).</p>
	Chemigation 2 pints/A	0.75 fl. oz.	<p>Apply to established and newly seeded grasses grown for seed or sod production before the boot stage. Apply through automated sprinkler irrigation systems with mechanical transfer loading system only. See MIXING, LOADING AND HANDLING INSTRUCTIONS section for complete details.</p> <p>Established grasses tolerant to BISON include bentgrasses, Kentucky bluegrass, fescues, ryegrasses, Bermuda grass, St. Augustine grass and zoysiagrass. Bison may also be used on seedling grasses such as Merion, Park, Delta, or common Kentucky bluegrasses, Pennlawn, chewings, Illahee, or Alta fescues, orchard grass, Highland, Seaside, or Astoria bentgrasses, perennial ryegrasses, Bahiagrass and zoysiagrass.</p>	

RESTRICTIONS AND PRECAUTIONS: Grasses grown for seed or sod production

- Do not allow livestock to graze in treated areas or feed treated grasses to livestock.
- Do not apply BISON to grasses grown for seed or sod production with backpack or hand-held equipment.

Distributed By:

Agriliance, LLC, P.O. Box 64089, St. Paul, MN 55164-0089

AgriSolutions and BISON™ are trademarks of Agriliance, LLC