

Bayer CropScience LP 800 N. Lindbergh Blvd. St. Louis MO, 63167 1-866-99BAYER (1-866-992-2937)

### **MOVENTO® HL**

For control of two-spotted spider mite in sugar beet.

EPA Reg. No. 264-1188

## FIFRA 2(ee) Recommendation

# FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF IDAHO, OREGON AND WASHINGTON

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its label.

### **CROP USE DIRECTIONS**

Apply specified dosage of MOVENTO HL early in the infestation as the population begins to develop or at early threshold for the target insect pest. Apply higher dosages specified within the crop specific sections when applied as a preventive application, for moderate to heavy insect pressure, or where longer residual control is desired. Degree of efficacy against labeled pests will be determined, in part, by the stage of pest development at application and infestation level of those pests.

Apply in adequate water for uniform coverage. For field crops, apply in a minimum of 10 GPA by ground and 5 GPA by aerial application. For vegetable and potato crops, apply in a minimum of 15 GPA by ground and 5 GPA by aerial application. MOVENTO HL may also be applied through overhead irrigation systems as designated in the CHEMIGATION section of the label under *Chemigation Application Instructions*.

MOVENTO HL must be tank mixed with a spray adjuvant / additive having spreading and penetrating properties to maximize leaf uptake and systemicity of the active ingredient within treated plants; please contact your local Bayer CropScience representative or PCA for specific recommendations by crop. The tank mixture of MOVENTO HL with an adjuvant / additive having sticking properties or crop protection product formulations containing built-in stickers have been shown to interfere with leaf uptake and should be avoided. Sufficient leaf tissue must be present for uptake and translocation of this product.

SUGAR BEET		
Pests Suppressed	Product Rate	
Two-spotted Spider Mite	(fl oz/A)	(lb ai/A)
	2.25 – 4.5	0.07 – 0.14
Foliar Application Restrictions:		
Pre-Harvest Interval (PHI): 28 day(s)		
Minimum interval between applications: 14 days		
Maximum MOVENTO HL allowed per crop season: 9 fl oz/A		
Maximum spirotetramat per crop season: 0.28 lb ai/A		

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577. For <u>PRODUCT USE</u> Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).