

 INSECTICIDE

PYRIPROXYFEN GROUP 7C INSECTICIDE

 **Skida™**  
35 WSP

**Insect Growth Regulator**

<b>ACTIVE INGREDIENT:</b>	<b>By Wt.</b>
*Pyriproxyfen .....	35.0%
<b>OTHER INGREDIENTS:</b> .....	65.0%
<b>TOTAL</b> .....	100.0%
*2-[1-methyl-2-(4-phenoxyphenoxy)ethoxy]pyridine	
Skida™ 35 WSP is a wettable powder formula containing 35% active ingredient.	

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See inside label booklet for First Aid, Precautionary Statements and Directions for Use.

FOR A MEDICAL EMERGENCY OR HELP WITH ANY SPILL, LEAK, FIRE OR EXPOSURE INVOLVING THIS MATERIAL, CALL DAY OR NIGHT, CHEMTREC 1-800-424-9300.

**EPA REG. NO. 34704-1173**

L090723-102425

MANUFACTURED FOR  
LOVELAND PRODUCTS, INC.®, P.O. BOX 1286, GREELEY, COLORADO 80632-1286

  
**Loveland**  
PRODUCTS.

## FIRST AID

<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency medical assistance, call **1-800-424-9300**. For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC **1-800-424-9300**.

For Chemical Emergency Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC Day or Night Within USA and Canada: **1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if inhaled, swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

### PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt,
- Long pants,
- Socks plus shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by disposing of equipment washwaters or rinsate. Avoid direct application and/or spray drift to bee hives.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### USE RESTRICTIONS ON ALL CROPS

- Apply this product only as specified in the EPA approved label and any supplemental labeling.
- **Do not apply this product through any type of irrigation system.**
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE), and Restricted Entry Interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants,
- Waterproof gloves,
- Socks plus shoes

## TANK MIXES

**NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of the user, applicator and/or applicator advisor, to the extent allowable by applicable law.**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## RESISTANCE MANAGEMENT

For resistance management, Skida 35 WSP is a Group 7C Insecticide. Any insect/mite population may contain individuals naturally resistant to Skida 35 WSP and other Group 7C insecticides/acaricides. The resistant individuals may dominate the insect/mite population if this group of insecticide/acaricides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay insecticide/acaricide resistance, take the following steps:

- Rotate the use of Skida 35 WSP or other Group 7C insecticides/acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides/acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticides/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Loveland Products, Inc. retailer, representative or call 1-800-424-9300. You can also contact your pesticide distributor or university extension specialist to report resistance.

Refer to specific crop sections for **"Managing Insect Resistance"** language applicable to that crop.

## WATER SOLUBLE PACKAGING

### MULTIPLE PACKAGING

This bag contains multiple water-soluble packets of Skida 35 WSP. Do not handle the packets with wet gloves or allow the packets to become wet prior to addition to spray tank. Do not break open packets. Refer to the **Table of Equivalents** to calculate the number of packets to use. If all packets are not used, close and reseal outer container to protect remaining packet(s).

RATES OF Skida 35 WSP PER ACRE	# OF WATER SOLUBLE PACKETS TO TREAT 4 ACRES
3 oz	3
4 oz	4
5 oz	5

## PRODUCT INFORMATION

**Skida 35 WSP is intended for use in Integrated Pest Management (IPM) or Insect Resistance Management (IRM) programs. Skida 35 WSP will not control insect adults, and it is recommended to be used in combination and/or rotation with other IPM or IRM materials. Contact your local state extension service for details.**

Skida 35 WSP acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Skida 35 WSP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Skida 35 WSP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides. Skida 35 WSP exhibits translaminar movement which means that Skida 35 WSP applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

### POME FRUIT (Crop Group 11-10)

Skida 35 WSP acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Skida 35 WSP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Skida 35 WSP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

*Contact'd*

**POME FRUIT (Crop Group 11-10) (Cont'd)**

CROPS	PESTS	PRODUCT RATE PER ACRE	INSTRUCTIONS
Apple Azarole Crabapple Loquat Mayhaw Medlar Quince Quince, Chinese Quine, Japanese Tejocote Cultivars, varieties, and/or hybrids of these	Codling Moth  Suppression of: Obliquebanded Leafroller Pandemis Leafroller	4 to 5 oz	Apply by ground with airblast equipment in a minimum of 50 gals/A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.  <b>1<sup>st</sup> Generation Codling Moth:</b> Apply at Biofix plus 100-degree days (this is usually close to petal fall). Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase Codling Moth activity of Skida 35 WSP. Follow local recommendations regarding in-season use and rates of oil in your area.  <b>Obliquebanded and Pandemis Leafroller:</b> Apply to the overwintering larvae at pink to petal fall tree growth stage. Skida 35 WSP does not directly control the larvae but inhibits normal pupation and adult emergence.
	Florida Red Scale Lecanium Scale San Jose Scale	4 to 5 oz	Apply by ground with airblast equipment in a minimum of 50 gals/A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.  Apply at delayed dormant through pink. For delayed dormant use a superior spray oil at the manufacturer's recommended rate of oil.  Under heavy scale pressure, use the higher rate of Skida 35 WSP.  <b>Scale:</b> For control in-season, make application in cover sprays when scale crawlers first emerge. The use of oil at the crawler emergence stage has been shown to improve control. Follow local recommendations regarding in-season use and rates of oil in your area.
	Green Apple Aphid Rosy Apple Aphid Spirea Aphid Spotted Tentiform Leafminer Western Tentiform Leafminer	3 to 5 oz	Apply by ground with airblast equipment in a minimum of 50 gals/A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.  <b>Leafminers:</b> Apply at peak moth flight for control of 1 <sup>st</sup> or 2 <sup>nd</sup> generation. This usually occurs at pre-pink to pink apple stage for 1 <sup>st</sup> generation and 2 <sup>nd</sup> cover for 2 <sup>nd</sup> generation.  <b>Rosy Apple Aphid:</b> Apply at ½" green apple growth stage.  <b>Green Apple &amp; Spirea Aphid:</b> Apply when aphids first appear, but before aphids are colonized. <b>TIMING IS CRITICAL FOR ACCEPTABLE APHID CONTROL.</b>

**POME FRUIT (Crop Group 11-10) (Cont'd)**

CROPS	PESTS	PRODUCT RATE PER ACRE	INSTRUCTIONS
Pear Pear, Asian	Pear Psylla	4 to 5 oz	<p>Apply by ground with airblast equipment in 75 to 400 gals/A with a superior spray oil at manufacturer's recommended rates for the stage of growth. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply a single application during the delayed dormant to pink stage of growth at 5 oz/A or make two applications from delayed dormant through petal fall stage of growth at 4 to 5 oz/A. Make application just before sustained egg lay is anticipated.</p> <p>For in-season control of pear psylla, use an insecticide recommended for this use.</p>
	Lecanium Scale San Jose Scale	4 to 5 oz	<p>Apply by ground with airblast equipment in 100 to 400 gals/A at delayed dormant with a superior spray oil at the manufacturer's recommended rates. Under heavy scale pressure, use the higher rate.</p> <p>Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>For control of scale in-season make application in cover sprays when scale crawlers first emerge. The use of oil at the crawler emergence stage has been shown to improve performance. Follow local recommendations regarding in-season use and rates of oil in your area.</p>
	Codling Moth  Suppression of: Obliquebanded Leafroller Pandemis Leafroller	4 to 5 oz	<p>Apply by ground with airblast equipment in 100 to 400 gals/A.</p> <p><b>1<sup>st</sup> Generation Codling Moth:</b> Apply at Biofix plus 100- degree days. Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase Codling Moth activity of Skida 35 WSP. Follow local recommendations regarding in-season use and rates of oil in your area.</p> <p>For additional generations apply other insecticides registered for Codling Moth control.</p>
<p><b>Note:</b> Skida 35 WSP may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p>			
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT SKIP ROWS DURING APPLICATION.</b></li> <li>• Do not apply within 45 days of harvest.</li> <li>• Do not make more than 2 Skida 35 WSP applications per growing season.</li> <li>• Do not exceed 10 oz of Skida 35 WSP per acre per season.</li> <li>• Do not apply earlier than 14 days after last Skida 35 WSP treatment.</li> <li>• Regardless of formulation, do not apply more than 0.218 lb pyriproxyfen per acre per season.</li> </ul>			

## SMALL FRUIT VINE CLIMBING CROPS (EXCEPT GRAPE)

(Subgroup 13-07E)

Not Registered for Use by New York

Skida 35 WSP acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Skida 35 WSP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Skida 35 WSP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PESTS	PRODUCT RATE PER ACRE	INSTRUCTIONS
Berry, Schisandra* Gooseberry* Grape, Amur River* Kiwifruit, Fuzzy Kiwifruit, Hardy Maypop* Cultivars, varieties, and/or hybrids of these  *Not Registered for Use by California	Scale Whitefly	5 oz	Apply with ground or air equipment as a full coverage spray (minimum of 5 gals/A by air or 50 gals/A by ground). Thorough coverage is critical for adequate control. It is essential to use adequate water volume to ensure thorough coverage.  Apply at either dormant, delayed dormant or an in-season cover spray. At dormant or delayed dormant use a spray oil at the manufacturer's recommended oil rate. In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area.
<p><b>Note:</b> Skida 35 WSP may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"><li>• Do not apply within 30 days of harvest.</li><li>• Do not make more than 2 Skida 35 WSP applications per growing season.</li><li>• Do not exceed 10 oz of Skida 35 WSP per acre per season.</li><li>• Do not apply earlier than 14 days after last Skida 35 WSP treatment.</li><li>• Regardless of formulation, do not apply more than 0.218 pyriproxyfen per acre per season.</li></ul>			

**STONE FRUIT**  
**(Crop Group 12)**

Skida 35 WSP acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Skida 35 WSP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Skida 35 WSP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PESTS	PRODUCT RATE PER ACRE	INSTRUCTIONS
Apricot Cherry Chickasaw Plum Damson Plum Fresh Prune Japanese Plum Nectarine Peach Plum Plumcot Prune	Lecanium Scale San Jose Scale  Suppression of: Peach Twig Borer Oriental Fruit Moth	4 to 5 oz	<p>Apply in 100 to 400 gals/A with ground equipment. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will ensure better coverage.</p> <p><b>Scale:</b> Apply at dormant, delayed dormant or as an in-season cover spray. At dormant, delayed dormant, use a spray oil at the manufacturer's recommended oil rate.</p> <p>Under heavy scale pressure, use the higher rate of Skida 35 WSP.</p> <p><b>Peach Twig Borer:</b> Skida 35 WSP has provided good suppression of Peach Twig Borer when applied delayed dormant through bloom sprays. Use a spray oil at the manufacturer's recommended rate for the stage of crop growth.</p>
<p><b>Note:</b> In-season sprays of Skida 35 WSP may not protect fruit from blemishes by scale.</p> <p>Skida 35 WSP may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply within 14 days of harvest.</li> <li>• Do not make more than 3 Skida 35 WSP applications per growing season.</li> <li>• Do not exceed 15 oz of Skida 35 WSP per acre per season.</li> <li>• Do not apply earlier than 14 days after last Skida 35 WSP treatment.</li> <li>• Regardless of formulation, do not apply more than 0.327 lb pyriproxyfen per acre per season.</li> </ul>			

**TREE NUTS**  
**(Crop Group 14)**

Skida 35 WSP acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Skida 35 WSP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Skida 35 WSP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PESTS	PRODUCT RATE PER ACRE	INSTRUCTIONS
Almond	Black Scale San Jose Scale  Suppression of: Peach Twig Borer	4 to 5 oz	Apply with ground equipment in 100 to 400 gals/A. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will ensure better coverage.  Apply at either dormant, delayed dormant or an in-season cover spray. At dormant, delayed dormant use a spray oil at the manufacturer's recommended oil rate. In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area.  Skida 35 WSP has provided good suppression of Peach Twig Borer when applied delayed dormant through bloom sprays.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Do not apply within 21 days of harvest.</li> <li>• Do not make more than 1 Skida 35 WSP application per growing season.</li> <li>• Do not exceed 5 oz of Skida 35 WSP per acre per season.</li> <li>• Regardless of formulation, do not apply more than 0.109 lb pyriproxyfen per acre per season.</li> </ul>			

*Cont'd*

**TREE NUTS (Crop Group 14) (Cont'd)**

CROPS	PESTS	PRODUCT RATE PER ACRE	INSTRUCTIONS
Beechnut Brazil Butternut Cashew Chestnut Chinquapin Filbert (Hazelnut) Hickory Macadamia Pecan Pistachio Walnut	Codling Moth Dusky Veined Aphid Walnut Aphid	5 oz	Apply with ground equipment in 100 to 400 gals/A. Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.  <b>1<sup>st</sup> Generation Codling Moth:</b> Apply at Biofix plus 100-degree days. Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase Codling Moth activity of Skida 35 WSP. Follow local recommendations regarding in-season use and rates of oil in your area.
	Black Scale Brown Soft Scale Italian Scale Lecanium Scale San Jose Scale Walnut Scale  Suppression of: Peach Twig Borer	4 to 5 oz	Apply with ground equipment in 100 to 400 gals/A at either dormant, delayed dormant or an in-season cover spray. The addition of a superior spray oil at dormant or delayed dormant may improve scale control. However, do not use a spray oil in oil sensitive crops such as Walnut. Follow manufacturer's recommendations regarding use and rates of oil for your area.  In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area.  Under heavy scale pressure, use the higher rate.  Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will ensure better coverage.
<p><b>Note:</b> Skida 35 WSP may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p>			
<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply within 21 days of harvest.</li> <li>• Do not make more than 2 Skida 35 WSP applications per growing season.</li> <li>• Do not exceed 10 oz of Skida 35 WSP per acre per season.</li> <li>• Do not apply earlier than 14 days after last Skida 35 WSP treatment.</li> <li>• Regardless of formulation do not apply more than 0.218 lb pyriproxyfen per acre per season.</li> </ul>			

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment.

**PESTICIDE STORAGE:** Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, and away from other pesticides, fertilizers, food or feed. Do not put concentrate or dilute into food or drink containers. Not for use or storage in or around the home.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable outer bag. Do not reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose of the empty bag in a sanitary landfill or by other procedures allowed by State and local authorities.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. To the extent consistent with applicable law, the buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS" AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESSED OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Skida 35 WSP is a trademark of Loveland Product, Inc.



## Insect Growth Regulator

## ACTIVE INGREDIENT:

*Pyriproxyfen	By Wt.	35.0%
OTHER INGREDIENTS:		65.0%
TOTAL		100.0%

\*2-[1-methyl-2-(4-phenoxyphenoxy)ethoxy]pyridine  
Skida™ 35 WSP is a wettable powder formula containing 35% active ingredient.

## KEEP OUT OF REACH OF CHILDREN CAUTION

## FIRST AID

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

## HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical assistance, call **1-800-424-9300**. For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC: **1-800-424-9300**.

MANUFACTURED FOR: LOVELAND PRODUCTS, INC.®, P.O. BOX 1286, GREELEY, COLORADO 80632-1286

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if inhaled, swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

## PESTICIDE STORAGE:

Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, and away from other pesticides, fertilizers, food or feed. Do not put concentrate or dilute into food or drink containers. Not for use or storage in or around the home.

## PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

## CONTAINER HANDLING

Nonrefillable outer bag. Do not reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose of the empty bag in a sanitary landfill or by other procedures allowed by State and local authorities.

See inside label booklet for additional Precautionary Statements and Directions for Use.

EPA REG. NO. 34704-1173

L090723-102425



↑  
PEEL HERE →

EPA EST. NO. 67545-AZ-001(G); 65387-AR-002(S)

First letter(s) in lot number correspond to letter(s) following the EPA Est. No.

NET WEIGHT: 2.5 lb (1.13 kg)

Each pouch 10 x 4 WT oz WSP.