

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1185A-S7 0117**

**Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.**

GROUP	9	15	HERBICIDES
-------	---	----	------------

### **Sequence® Herbicide**

**This supplemental label expires on 01/05/2020 and must not be used or distributed after this date.**

#### **Active Ingredients:**

*Glyphosate: N-(phosphonomethyl) glycine .....	21.8%
**S-metolachlor (CAS No. 87392-12-9).....	29.0%
Other Ingredients:	49.2%
<b>Total:</b>	<b>100.0%</b>

\*Contains 2.25 pounds of glyphosate acid per U.S. gallon.

\*\*Contains 3 pounds of S-metolachlor per U.S. gallon.

Sequence® Herbicide is formulated as an emulsion in water (EW).

**KEEP OUT OF REACH OF CHILDREN.**

## **CAUTION**

**EPA Reg. No. 100-1185**

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Sequence Herbicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## **DIRECTIONS FOR USE**

### **Postemergence Over-the-Top Applications for Roundup Ready® Flex Cotton Only (For Use In AL, AR, AZ, CA, FL, GA, KS, KY, LA, MD, MO, MS, NC, NM, OK, SC, TN, TX, and VA Only)**

**When to Apply:** Postemergence, post-directed or hooded sprayer applications in Roundup Ready Flex cotton only.

Follow directions in the **APPLICATION AND MIXING PROCEDURES, SPRAY ADDITIVES**, and **APPLICATION EQUIPMENT AND METHODS** sections. Refer to **WEED CONTROL** section (Tables 1-3), for weeds controlled and application rates.

### **Use Precautions for Postemergence Over-the-Top Use on Roundup Ready Flex Cotton**

- Crop canopy interference can reduce spray coverage on target weeds and soil and hinder weed control. In large cotton, to improve spray coverage of target weeds, apply Sequence Herbicide in 12 or more gallons of water per acre.

### **Use Restrictions for Postemergence Over-the-Top Use on Roundup Ready Flex Cotton**

- Make postemergence applications from cotyledon stage to the 10-leaf stage (not to exceed 12 inches tall) of cotton development. Do not apply later as severe injury, including yield loss, could occur.
- Do not exceed 2.5 pt (0.94 lb S-metolachlor and 0.7 lb glyphosate) of Sequence Herbicide per acre in a single application on cotton with less than 5 leaves.
- Apply up to 3.5 pt of Sequence Herbicide per acre in a single application from the 5-leaf through the 10-leaf stage of cotton.
- Do not exceed 3.5 pt (1.3 lb S-metolachlor) of Sequence Herbicide per acre per season applied postemergence.
- Use only water as the carrier for postemergence and post-directed applications in Roundup Ready Flex cotton.
- Do not use Sequence Herbicide postemergence if cotton plants are under stress, including, but not limited to, that caused by drought, insect, disease, or injury from cultivation. Cotton leaves may exhibit necrotic spotting that will not affect normal plant development or cotton yield.

- Do not harvest within 100 days of postemergence over-the-top application of Sequence Herbicide.
- Do not include AMS or other adjuvants when applications are made postemergence to the cotton.
- Do not graze or feed forage or fodder from cotton to livestock.
- Do not use in Gaines County, TX.
- Do not apply to Taloka silt loam.
- Do not apply Sequence Herbicide to emerged **conventional** cotton, as severe crop injury will occur.
- Do not exceed a total of 1.9 lb active ingredient of S-metolachlor per acre per calendar year on coarse textured soils or 2.47 lb active ingredient of S-metolachlor per acre per year on medium or fine-textured soils resulting from all applications of Sequence Herbicide, Dual Magnum, Dual II Magnum or other S-metolachlor containing products.

Sequence® Herbicide and the Syngenta logo are Trademarks of a Syngenta Group Company

Roundup Ready® is a registered Trademark of Monsanto Corporation

©2017 Syngenta