

FINISH™

Guaranteed Analysis
2-14-3

Total Nitrogen (N)	2.00%
1% Urea Nitrogen	
1% Nitrate Nitrogen	
Available Phosphate (P ₂ O ₅)	14.0%
Soluble Potash (K ₂ O)	3.0%
Molybdenum (Mo)	2.0%

Derived from urea, phosphoric acid, potassium hydroxide, potassium nitrate and sodium molybdate.

CAUTION

Keep out of Reach of Children • May Cause Eye Irritation

USER SAFETY RECOMMENDATIONS

Avoid contact with skin, eyes or clothing. Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove wet clothing immediately. Wash skin thoroughly and put on clean clothing.

FIRST AID

Call a poison control center or doctor for treatment advice. Have product container or label with you when calling a poison control center, or doctor, or going for treatment. If in eyes: Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. If on skin: Take off contaminated clothing and rinse skin immediately with plenty of water for 15 to 20 minutes.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

WARNING: Only use this product on soils that respond to molybdenum. Crops high in molybdenum are toxic to grazing animals (ruminants).

STORAGE AND DISPOSAL

Store this product out of direct sunlight, in a cool place above 40°F. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Dispose of empty container in a sanitary landfill or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals>.

NET CONTENTS:

Net Weight:10.35 lbs/gallon at 68°F

□2.5 gal. (25.5 lb) □55 gal. (569.25 lb) □10 liter (12.22 kg) □208 liter (254.18 kg)
□5.0 gal. (51 lb) □275 gal. (2846.25 lb) □20 liter (24.44 kg) □1040 liter (1270.88 kg)

FINISH™

F001601



Z-FINISH

Manufactured by Stoller Enterprises, Inc.
4001 W. Sam Houston Parkway North, Suite 100 | Houston Texas 77043 USA
Phone: 1-800-539-5283 or 713-461-1493 | Fax: 713-461-4467

GENERAL INFORMATION

Use Finish as a foliar application on any grain, fruit, or storage tissue crop in preparation for harvest. During the ripening process the cells of the seed, fruit, or storage tissue changepolarity and generate exothermic energy, which results in the loss of carbohydrates and proteins out of the harvestable tissue and back into the mother plant due to an additional energy demand for survival during its late period in life. Finish will disable this process and force all of the sugars and proteins to remain in the grain, fruit or storage tissue resulting in better sizing and fill.

DIRECTIONS FOR USE

It may be in violation of state law to use this product inconsistent with its labeling; always read and follow the label when using this product.

Use Finish after cell sizing is complete and moisture loss has commenced to prepare the seed, fruit, or storage tissue for harvest and increase dormancy for better shelf life or storage properties. Apply the recommended amount in a volume sufficient to provide thorough coverage.

RECOMMENDED USES

Apply Finish at 1 quart per acre during senescence or 2 quarts per acre if the crop remains vigorous on the following Field Crops: alfalfa, asparagus, bean crops, canola, clover, corn, cotton, cruciferous crops, cucurbits, milo, melons, onion/garlic, peanut, pepper, potato, rape (canola), rice, soybean, sugar beet, sugarcane, sunflower, tobacco, tomato, vegetable crops, wheat/barley/oat; or Fruit and Nut Crops: almond, apple, apricot, avocado, banana, berry crops, cherry, citrus, coffee, fig, filbert, grapes, hazelnut, oil palm, olive, persimmon, peach, pear, pecan, pistachio, plums/prunes, pomegranate, strawberry and walnut.

APPLICATION TIMING

Apply Finish 20 days before harvest unless conditions dictate otherwise. On corn, apply Finish at the recommended rate at R5. On soybean, apply Finish when the upper pod has well developed bean seeds. On wheat and other small grains, apply Finish immediately during the advanced dough stage. On Fruit and nut trees, apply Finish at color break.

Notice: The information contained herein is true and accurate to the best of our knowledge and belief, but is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturer nor the seller make any warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instruction. Read and follow the label directions carefully. **CONDITIONS OF SALE:** 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond sellers control, Sellers liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.