

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

THIAMETHOXAM	GROUP	4A	INSECTICIDE
CYANTRANILIPROLE	GROUP	28	INSECTICIDE



Insecticide

For foliar and systemic control of listed insect pests in ornamental plants, residential and commercial landscapes, fruit and nut trees (non-bearing), Christmas tree seedlings (liner production and field application) and listed vegetable transplants grown for sale to consumers.

Active Ingredient:

Thiamethoxam ¹	20.0%
Cyantraniliprole ²	20.0%

Other Ingredients: 60.0%

Total: 100.0%

¹ CAS No. 153719-23-4

² CAS No. 736994-63-1

Mainspring® Xtra is a water-dispersible granule that contains 3.2 ounces of thiamethoxam and 3.2 ounces of cyantraniliprole per pound of formulated product.

¹ Thiamethoxam belongs to the neonicotinoids chemical class

² Cyantraniliprole belongs to the anthranilic diamide chemical class

**KEEP OUT OF REACH OF CHILDREN./
MANTENGASE FUERA DEL ALCANCE DE
LOS NIÑOS.**

CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet. Vea mas declaraciones de precaucion e instrucciones del uso en folleto.

EPA Reg. No. 100-1424

EPA Est. 67545-AZ-1

SCP 1424B-L1A 0225

48 oz

Net Weight

Non-refillable container

syngenta®

FIRST AID

If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOTLINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or
Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),
Call
1-800-888-8372

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton™ ≥14 mils
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates, oysters and shrimp.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds and may cause possible effects to pollinators from exposure to translocated residues in blooming crops. Do not apply this product or allow it to drift to blooming crops while bees are foraging in/or adjacent to the treatment area.

Surface Water Advisory

This product may impact surface water quality due to spray drift and runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground-water. This product is classified as having high potential for reaching surface water via runoff for several months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of thiamethoxam water from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

(See manual at the following internet address:
<http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>).

continued...

PRECAUTIONARY STATEMENTS (continued)

Surface Water Protection Statements

- For foliar applications: Do not apply during rain.
- **DO NOT** cultivate within 30 feet of the aquatic area to allow growth of a vegetative filter strip.
- **DO NOT** apply within 50 ft of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, wetlands or natural ponds, estuaries, and commercial fish farm ponds).

Groundwater Advisory

This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Spray Drift Advisory

Mainspring Xtra insecticide may be applied by ground application equipment.

Physical or Chemical Hazards

DO NOT use with or store near oxidizing agents since a hazardous chemical reaction may occur.



PROTECTION OF POLLINATORS

APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.

Look for the bee hazard icon  in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to:

www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER**

FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and of Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

See individual use sites for specific pollinator protection application restrictions. If none exist, follow these directions for foliar applications to commercially-grown plants and ornamentals that are attractive to pollinators and non-agricultural use sites:

FOR FOOD/FEED CROPS AND COMMERCIAL GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS

Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset
- The application is made to the target site when temperatures are below 55°F
- The application is made in accordance with a government-initiated public health response
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48 hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify beekeepers no less than 48 hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.

FOR NON-AGRICULTURAL USES

- Do not apply Mainspring Xtra while bees are foraging. Do not apply Mainspring Xtra to plants that are flowering. Only apply after all flower petals have fallen off.

ENDANGERED AND THREATENED SPECIES PROTECTION REQUIREMENTS

Before using this product, you must obtain any applicable Endangered Species Protection Bulletins ('Bulletins') within six months prior to or on the day of application. To obtain Bulletins, go to Bulletins Live! Two (BLT) at <https://www.epa.gov/pesticides/bulletins>. When using this product, you must follow all directions and restrictions contained in any applicable Bulletin(s) for the area where you are applying the product, including any restrictions on application timing if applicable. It is a violation of federal law to use this product in a manner inconsistent with its labeling, including this labeling instruction to follow all directions and restrictions contained in any applicable Bulletin(s). For general questions or technical help, call 1-844-447-3813, or email ESPP@epa.gov.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours except for automatic fogger use in greenhouses (See "Automatic Cold Fogger Use in Greenhouses" below). PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

Automatic Cold Fogger Use in Greenhouses: Early entry is based on ventilation criteria.

- If there is no ventilation, the REI is 24 hours.
- Early entry is permitted (with the PPE listed above) and respiratory protection consisting of a protection factor (PF) 10 respirator fitted with an organic-vapor removing filter) after four (4) hours to perform short duration activities such as opening windows or turning on fans or ventilation systems.
- Early entry is permitted (with the PPE listed above) after two (2) hours of ventilation using fans or other mechanical ventilation systems.
- Early entry is permitted (with the PPE listed above) after four (4) hours of ventilation using vents, windows or other passive ventilation.
- Early entry is permitted (with the PPE listed above) after ten (10) complete air exchanges have been completed.

After 12 hours, if the greenhouse has been ventilated according to the above criteria, no PPE is required to enter the treated area.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Professional applications to golf courses, residential, industrial and commercial lawns and landscapes, and sports fields are not within the scope of the Worker Protection Standard. Do not enter or allow others to enter the treated area until sprays have dried.

Do not enter treated areas without protective clothing until sprays have dried.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY AND/OR POOR INSECT CONTROL.

USE INFORMATION

RESISTANCE MANAGEMENT

THIAMETHOXAM	GROUP	4A	INSECTICIDE
CYANTRANILPROLE	GROUP	28	INSECTICIDE

Some insect pests are known to develop resistance to products after repeated use. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area. Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects on this label.

Mainspring Xtra contains a Group 4 insecticide (thiamethoxam, belonging to the neonicotinoid class of chemistry) and a Group 28 insecticide (cyantranilprole, belonging to the diamide class of chemistry). Insect biotypes with acquired or inherent resistance to Group 4 or Group 28 insecticides may eventually dominate the insect population if Group 4 or Group 28 insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by Mainspring Xtra or other Group 4 or Group 28 insecticides.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

In order to maintain susceptibility to these classes of chemistry:

- Avoid using Group 4 and/or Group 28 insecticides exclusively for season-long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, apply Mainspring Xtra or other Group 4 and/or Group 28 insecticides using a "treatment window" approach. A treatment window is a period of time as defined by the stage of crop development and/or the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, there may either be single or consecutive applications (seed treatment, soil, foliar, unless otherwise stated in the Directions for Use) of the Group 4 and/or Group 28 insecticides. Do not exceed the maximum Mainspring Xtra allowed per growing season.
- Following a treatment window of Group 4 and/or Group 28 insecticides, rotate to a treatment window of effective products with a different mode of action before making additional applications of Group 4 and/or Group 28 insecticides.

- A treatment window rotation, along with other IPM practices for the crop and use area, is considered an effective strategy for preventing or delaying a pest's ability to develop resistance to these classes of chemistry.
- If resistance is suspected, do not reapply Mainspring Xtra or other Group 4 or Group 28 insecticides.

Other Insect Resistance Management (IRM) practices include:

- Incorporating IPM techniques into your insect control program.
- Monitoring treated insect populations for loss of field efficacy.
- Using tank-mixtures or premixes with insecticides from a different target site of action group as long as the involved products are all registered for the same crop outlet and effective rates are applied.

For additional information on Insect Resistance Management:

- Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

APPLICATION EQUIPMENT CLEANING

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all application equipment to reduce the risk of forming hardened deposits that might become difficult to remove. Drain application equipment. Thoroughly rinse application equipment and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. **DO NOT** clean equipment near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

CALIBRATION OF APPLICATION EQUIPMENT

Proper maintenance and calibration of spraying and chemigation equipment are essential for optimal insect pest control. If you have questions about calibration, contact a State Extension Service specialist, the equipment manufacturer or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

MIXING PROCEDURES

Prepare only the amount of spray mixture that is needed for the immediate application. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation.

Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

Mainspring Xtra Applied Alone

Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the Mainspring Xtra to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the Mainspring Xtra has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Mainspring Xtra + Tank Mixtures

If using Mainspring Xtra in a tank mixture, it is the user's responsibility to observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. Do not exceed any label dosage rate and follow the most restrictive directions for use and precautionary language of each product in the mixture (for example, first aid from one product, spray drift management from another). Do not mix this product with any product, which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add the tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) such as Mainspring Xtra, liquid flowables, liquids and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

Important: When using Mainspring Xtra in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including Mainspring Xtra. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

When an adjuvant is to be used with this product, use an adjuvant that meets the standard of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program. Mainspring Xtra is compatible with most commonly used pesticides, crop oils, adjuvants, and nutritional sprays. However, since it is not possible to test all possible mixtures, the user should pre-test to assure the physical compatibility and lack of phytotoxic effect of any proposed mixtures with Mainspring Xtra.

Tank Mix Compatibility

Mainspring Xtra is compatible with most insecticide, fungicide, and foliar nutrients products. However, test the physical compatibility of Mainspring Xtra with tank-mix partners before use. To determine the physical compatibility of Mainspring Xtra with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Spray Drift Management

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions. For broadcast applications made at planting or prior to the emergence of crops, applicators are required to use a coarse or coarser droplet size (ASABE S572.1). For all other broadcast applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce the effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be

indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. Do not apply when wind speeds exceed 10 mph at the application site. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

POLLINATOR PRECAUTIONS

- Mainspring Xtra is highly toxic to bees exposed to direct treatment on blooming crops or weeds and may cause possible effects to pollinators from exposure to translocated residues in blooming crops.
- Do not apply Mainspring Xtra or allow it to drift to blooming crops while bees are foraging in or adjacent to the treatment area. This is especially critical if there are adjacent orchards that are blooming (Refer to **Spray Drift Precautions** for additional information).
- If bees are foraging in the ground cover and it contains any blooming plants or weeds, always remove flowers before making an application. This may be accomplished by mowing, disking, mulching, flailing, or applying a labeled herbicide.
- Consult with your local cooperative extension service or state agency responsible for regulating pesticide use for additional pollinator safety practices.

USE INFORMATION – ORNAMENTAL LANDSCAPE PLANTS

Mainspring Xtra is a broad-spectrum insecticide that is effective when applied at label rates to the foliage. Mainspring Xtra is active against listed sucking and chewing insect pests by contact and ingestion.

Application to Ornamental Plants

Mainspring Xtra may be used on most ornamental plants in residential landscaped areas and landscaped areas around institutional, public, commercial and industrial buildings; parks; recreational areas; and athletic fields (including trees, shrubs, flowers, evergreens, foliage plants, and groundcovers). Test Mainspring Xtra alone or the mixture to be applied (including adjuvants) on a small number of plants to determine any adverse plant safety effects before making large-scale applications.

Applications of Mainspring Xtra to yellow varieties of Honey Locust (*Gleditsia triacanthos*) may result in leaf chlorosis and leaf abscission. The effects are temporary. Test Mainspring Xtra alone or the mixture to be applied on a small number of the Honey Locust varieties you intend to treat and determine any adverse plant safety effects before making large-scale applications.

- DO NOT** apply this product, by any application method, to linden, basswood, or *Tilia* species.

APPLICATION PROCEDURES

Foliar Applications

Make foliar applications in an adequate water volume to achieve thorough and uniform coverage without excessive runoff (to drip). Use of an adjuvant applied with Mainspring Xtra may increase efficacy. Select an adjuvant that does not bind Mainspring Xtra to the leaf surface, limiting absorption into the foliage. Mainspring Xtra can be applied in a range of spray volumes (concentrate to dilute spray volumes), provided thorough and uniform coverage is obtained.

Handheld Application Equipment

Applications can be made through most types of low-pressure handheld application equipment.

DO NOT apply through any type of handheld fogger equipment.

Application Timing

Begin foliar applications when labeled pests first appear or when economic thresholds have been reached.

Apply Mainspring Xtra preventatively to plants that have low economic thresholds or are prone to infestation by listed insect pests.

Spray Drift Precautions

As with all crop protection products, it is important to avoid off-target movement. Do not allow spray to drift onto adjacent land, crops, or aquatic areas. To avoid spray drift:

- Make applications when wind velocity favors on-target product deposition (approximately 3-10 mph). Do not apply when wind velocity exceeds 10 mph. Do not make applications when wind gusts approach 10 mph.
- Do not make applications when wind direction is toward the aquatic area to reduce the risk of exposure to sensitive aquatic areas.
- Do not make applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with increased height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.
- Use the largest droplet size consistent with good pest control. Small droplets are more prone to spray drift, and can be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- For broadcast applications made at planting or prior to the emergence of crops, applicators are required to use a coarse or coarser droplet size (ASABE S572.1). For all other broadcast applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Apply as close to target plants as practical to obtain a good spray pattern for adequate coverage.

SPECIFIC USE DIRECTIONS – ORNAMENTAL LANDSCAPE PLANTS

Foliar Applications

Pests	Rate	Use Restrictions
Ants (excluding harvester, carpenter and Pharaoh ants) Aphids Beet Armyworm Cabbage Looper Cabbage Webworm Corn Earworm Diamondback Moth Fall Armyworm Flea Beetles Imported Cabbageworm Southern Armyworm Sugarbeet Armyworm Yellowstriped Armyworm	5-10.0 oz/A or 0.6-1.1 oz/5,000 sq ft	Make the first application when economic action levels for the target pest are reached. For crops without action levels, make the first application before pest populations cause significant damage to the host plant. A second application, if needed, may be made but no sooner than 14 days. Apply higher use rates in rate range for longer control or high insect populations. For leaf-gall-forming insects, apply during egg laying of the generation being targeted.
Adelgids (including Hemlock Woolly Adelgid) Lace Bugs Leaf Beetles Leafhoppers Leafminers Leaf gall forming insects Mealybugs Midges Plant Bug Psyllids (including Asian Citrus Psyllid) Sawflies Soft Scales Thrips – foliar feeding Whiteflies	10.5 oz/A or 1.2 oz/5,000 sq ft	For concentrated spray volumes, apply the same amount of product per unit area as would be applied with dilute spray volumes.

- DO NOT** exceed 21 oz/A of product (0.2625 lb of thiamethoxam/A and 0.2625 lb of cyantraniliprole/A) per year.
- DO NOT** apply more than 0.2625 lb ai/A thiamethoxam or 0.4 lb ai/A cyantraniliprole per calendar year regardless of formulation or method of application.
- For specific information about developmental stages of the target pest, associated damage and action thresholds to properly time applications, consult with your State Cooperative Extension Service.

USE INFORMATION – ORNAMENTAL PLANTS, NON-BEARING FRUIT AND NUT TREES (DEFINED AS CROPS THAT WILL NOT BEAR FRUIT WITHIN ONE YEAR OF APPLICATION), CHRISTMAS TREE SEEDLINGS (LINER PRODUCTION AND FIELD APPLICATION), AND LISTED VEGETABLE TRANSPLANTS GROWN FOR SALE TO CONSUMERS

Mainspring Xtra is a broad-spectrum insecticide that is effective when applied at label rates to the foliage, soil and growing media. Mainspring Xtra is active against listed sucking and chewing insect pests by contact and ingestion.

Application to Ornamental Plants

Mainspring Xtra can be used on most ornamental plants. Test Mainspring Xtra alone or the mixture (including adjuvants) to be applied on a small number of plants to determine any adverse plant safety effects before making large-scale applications.

Among container-grown ornamentals and ornamentals grown in planting or liner beds, some plant species may be sensitive to fertilizers. Determine the plant safety of Mainspring Xtra applied with fertilizers. Test the mixture on a small number of plants to determine any adverse plant safety effects before making large-scale applications.

Applications of Mainspring Xtra to yellow varieties of Honey Locust (*Gleditsia triacanthos*) may result in leaf chlorosis and leaf abscission. The effects are temporary. Test Mainspring Xtra alone or the mixture to be applied on a small number of the Honey Locust varieties you intend to treat and determine any adverse plant safety effects before making large-scale applications.

APPLICATION PROCEDURES

Foliar Applications

Make foliar applications in an adequate water volume to achieve thorough and uniform coverage without excessive runoff (to drip). Use of an adjuvant applied with Mainspring Xtra may increase efficacy. Select an adjuvant that does not bind Mainspring Xtra to the leaf surface, limiting absorption into the foliage. Mainspring Xtra can be applied in a range of spray volumes (concentrate to dilute spray volumes), provided thorough and uniform coverage is obtained.

Handheld Application Equipment

Applications can be made through most types of low-pressure handheld application equipment.

DO NOT apply through any type of handheld fogger equipment.

DO NOT apply with aerial equipment.

Automatic Cold Fogger Applications – Greenhouses

Applications can be made in greenhouses with automatic cold fogger equipment. Apply the same amount of Mainspring Xtra per unit area as would be applied in a dilute spray volume over the same area.

Do not apply through fogger equipment requiring workers to be present in the greenhouse during the fogger application.

Important: Read and follow the REI and PPE requirements for **Automatic Cold Fogger Use in Greenhouses** in the Worker Protection Standard **AGRICULTURAL USE REQUIREMENTS** box.

Soil or Soil-less Media Applications

Mainspring Xtra is effective as a soil and systemic insecticide when applied to the soil using band, broadcast, or drench applications. When applying Mainspring Xtra as a drench to plants in pots and containers, apply enough solution to sufficiently wet $\frac{1}{2}$ - $\frac{3}{4}$ of the root zone without allowing runoff out of the container. With irrigation or rainfall, move Mainspring Xtra into the root zone of plants or the location of soil-dwelling pests, preferably within 3 days of treatment and, optimally, within 24 hours. Excessive watering or heavy rainfall may reduce the effectiveness of Mainspring Xtra. Allow a minimum of 7 days after watering in for maximum uptake of Mainspring Xtra by the plant's root system.

Do not apply to soils or growing media above field or pot capacity. Do not apply to waterlogged soil or growing media.

Application Timing

Begin foliar applications when labeled pests first appear or when economic thresholds have been reached.

For applications made to growing media (soil or soil-less media), Mainspring Xtra must move to the feeding site of the target pest, whether this is in the root zone or the above-ground portion of the host plant. Therefore, control of pests from the systemic activity of Mainspring Xtra may be delayed for 1 or more weeks depending on plant size, rate of uptake from the soil, or rate of translocation within the plant, etc.

Systemic activity in large plants such as those in field nurseries may require several weeks for control of stem and foliar pests. Apply Mainspring Xtra preventatively to plants that have low economic thresholds or are prone to infestation by listed insect pests.

Application through Irrigation Systems

Apply Mainspring Xtra only through micro-irrigation systems (for example, drip, trickle, spaghetti tubes, and micro-sprinklers), overhead irrigation, ebb-and-flow and flood-floor irrigation, or hand-held or motorized calibrated irrigation equipment. Do not apply this product through any other type of irrigation equipment. Do not overwater for a minimum of 7 days after watering in for maximum uptake of Mainspring Xtra by the plant's root system. See additional precautions regarding irrigation system application below.

Chemigation

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Plant injury, lack of effectiveness, or illegal pesticide residues in the plant may result from non-uniform distribution of treated water.

The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or Venturi injector) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Public Water Systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or Venturi injector) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Ebb-and-Flow and Flood-Floor Application

To determine the water volume uptake of Mainspring Xtra in ebb-and-flow and flood-floor applications, bring a minimum of 10 plants up to a known field capacity and allow them to dry out for one to two days. Re-wet these plants to determine how much water on average each plant will absorb to bring it back to field capacity. Use the water volume absorbed per plant, multiply by the number of pots of the same size being treated and add the minimum water volume to irrigate the area. This should minimize the return back to the storage tank. Re-use the returned volume with subsequent irrigation or nutrients on the same plants. Dilute Mainspring Xtra in a minimum of 100 gallons for ebb-and-flow and flood-floor application.

Spray Drift Precautions

As with all crop protection products, it is important to avoid off-target movement. Do not allow spray to drift onto adjacent land, crops, or aquatic areas. To avoid spray drift:

- Make applications when wind velocity favors on-target product deposition (approximately 3-10 mph). Do not apply when wind velocity exceeds 10 mph. Do not make applications when wind gusts approach 10 mph.
- Do not make applications when wind direction is toward the aquatic area to reduce the risk of exposure to sensitive aquatic areas.
- Do not cultivate or plant crops within 25 ft of the aquatic area as to allow growth of a vegetative filter strip.
- Do not make applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with increased height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.
- Use the largest droplet size consistent with good pest control. Small droplets are more prone to spray drift, and can be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- For broadcast applications made at planting or prior to the emergence of crops, applicators are required to use a coarse or coarser droplet size (ASABE S572.1). For all other broadcast applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Apply as close to target plants as practical to obtain a good spray pattern for adequate coverage. Do not make applications more than 10 ft above the crop canopy.

SPECIFIC USE DIRECTIONS

Foliar Applications

Ornamental plants, non-bearing fruit and nut trees (defined as crops that will not bear fruit within one year of application), and Christmas tree seedlings grown in greenhouses, lath and shade houses, containers, field nurseries and interiorscapes

Pests	Rate	Use Restrictions
Ants (excluding harvester, carpenter and Pharaoh ants) Aphids Beet Armyworm Cabbage Looper Cabbage Webworm Corn Earworm Diamondback Moth Fall Armyworm Flea Beetles Imported Cabbageworm Southern Armyworm Sugarcane Armyworm Yellowstriped Armyworm	2.5-5.0 oz/100 gal or 5-10.0 oz/A or 0.6-1.1 oz 5,000 sq ft	Make the first application when economic action levels for the target pest are reached. For crops without action levels, make the first application before pest populations cause significant damage to the host plant. Re-apply as needed but no sooner than every 14 days. Apply higher use rates in rate range for longer control or high insect populations. For leaf- gall-forming insects, apply during egg laying of the generation being targeted.
Adelgids (including Hemlock Woolly Adelgid) Lace Bugs Leaf Beetles Leafhoppers Leafminers Leaf gall forming insects Mealybugs Midges Plant Bug Psyllids (including Asian Citrus Psyllid) Sawflies Soft Scales Thrips – foliar feeding Whiteflies	10.5 oz/100 gal or 10.5 oz/A or 1.2 oz/5,000 sq ft	For concentrated spray volumes, apply the same amount of product per unit area as would be applied with dilute spray volumes.

- For ornamental crops grown outdoors, do not exceed 21 oz/A of product (0.2625 lb of thiamethoxam/A and 0.2625 lb of cyantraniliprole/A) per year.
- For ornamental crops grown indoors, do not exceed 21 oz/A of product (0.2625 lb of thiamethoxam/A and 0.2625 lb of cyantraniliprole/A) per crop.
- Do not apply more than 0.2625 lb ai/A thiamethoxam or 0.4 lb ai/A cyantraniliprole per calendar year regardless of formulation or method of application.
- For specific information about developmental stages of the target pest, associated damage and action thresholds to properly time applications, consult your State Cooperative Extension Service.

Applications to Soil or Other Growing Media

Ornamental plants, non-bearing fruit and nut trees (defined as crops that will not bear fruit within one year of application) and Christmas tree seedlings grown in greenhouses, lath and shade houses, containers, field nurseries and interiorscapes

Pests	Rate	Use Restrictions
Aphids Fungus gnats Leaf Beetles Leafminers Mealybugs Root Aphid Root Weevil larvae (including <i>Diaprepes abbreviatus</i>) Soft Scales Thrips – foliar feeding (suppression) Whiteflies	5-10.5 oz/100 gal or 10.5 oz/A or 1.2 oz/5,000 sq ft	Apply to the soil or growing media to help prevent pests in situations where tolerance of insects or damage is low. Allow a minimum of 1 week for smaller plants and at least 2 weeks for large plants (typically found in nurseries, plantations) to translocate Mainspring Xtra to feeding sites of the target pest. Re-apply as needed, but no sooner than every 14 days. Apply as a broadcast, band, or drench.
Larvae (Grubs) of: <i>Aphodius</i> spp. Asiatic garden beetle Billbugs European chafer Green June beetle Japanese beetle May or June beetle Northern masked chafer Oriental beetle Scarab beetle Southern masked chafer	7.5-10.5 oz/100 gal	Apply from adult flight through peak egg hatch of the targeted species. Apply as a drench treatment to containers and over the active root zone. For drench and chemigation applications made to containerized plant material, apply sufficient volume to moisten the media around the root zone without loss from the bottom of the container.
Fire Ants	1.3-3.8 oz/10 gal	Individual Mound Treatment: For small mounds (<6 inch in diameter), apply 1 gal of water per mound and for larger mounds apply 2-3 gal of dilute solution per mound for optimum control. Direct the application to the center of the mound, including a 6-inch diameter circle.

- For broadcast applications, typical water volume is 1.5 - 5 gal/1,000 sq ft
- For ornamental crops grown outdoors, do not exceed 21 oz/A of product (0.2625 lb of thiamethoxam/A and 0.2625 lb of cyantraniliprole/A) per year.

- For ornamental crops grown indoors, do not exceed 21 oz/A of product (0.2625 lb of thiamethoxam/A and 0.2625 lb of cyantraniliprole/A) per crop.
- Do not apply more than 0.2625 lb ai/A thiamethoxam or 0.4 lb ai/A cyantraniliprole per calendar year regardless of formulation or method of application.
- Do not over-water for a minimum of 7 days after watering in for maximum uptake of Mainspring Xtra by the plant root system.
- For specific information about developmental stages of the target pest, associated damage, and action thresholds to properly time applications, consult with your State Cooperative Extension Service.

Banded Application

For banded applications, apply uniformly over the row or active root zone in a minimum water volume of 5 gallons per 1,000 linear feet. Irrigate within 24 hours of application if sufficient rainfall (1/4-1 inch of water) doesn't occur.

Mainspring Xtra rate (ounces of product) per 1,000 linear feet of row for specific row spacing

Rate/A	Lb ai/A	Row Spacing											
		3 ft	4 ft	4.5 ft	5 ft	6 ft	7 ft	8 ft	9 ft	10 ft	11 ft	12 ft	
10.5 oz/A	0.26	0.7	1.0	1.1	1.2	1.4	1.7	1.9	2.2	2.41	2.66	2.9	

Drench Application

When applying Mainspring Xtra as a drench, apply enough solution to sufficiently wet 1/2 - 3/4 of the root zone without allowing runoff out of the container. Use the following chart as a guideline for the number of pots that this volume will treat based on the size of the pot and drench volume. Use higher drench volumes for woody ornamentals, containers having multiple plants, and those crops with longer production times or needing longer protection.

Note: Drench volumes will vary based on the porosity of the growing media. Conduct your own tests to determine correct drench volumes for your growing media.

Pot Size	Suggested Drench Volume fl oz/pot	Pots Treated per 100 gal
4-inch standard	2-3	6,400-4,267
6-inch standard	3-4	4,267-3,200
8-inch standard	5-10	2,560-1,280
10-inch standard	12.5-25	1,024-512

Foliar Application to Vegetable Plants Grown for Sale to Consumers

For tomatoes, peppers, eggplant, and cucurbit plants grown for sale to consumers

Crop	Pests	Rate	Use Restrictions
Fruiting Vegetables Eggplant Ground-cherry Pepino Peppers (bell, chili, cooking, pimiento, and sweet) Tomatillo Tomato	Aphids Beet Armyworm Colorado Potato Beetle Fall Armyworm Flea Beetles Hornworm species Leafhoppers Leafminers Loopers Potato Psyllid Southern Armyworm Thrips (Foliar Feeding) Tomato Fruitworm Tomato Pinworm Whiteflies Yellowstriped Armyworm	5.0–7.0 oz/A or 0.6–0.8 oz/ 5,000 sq ft	Apply before pests reach damaging levels. Apply higher rates within the listed rate range for heavy infestations.

- For transplants grown outdoors, do not exceed 14.0 oz/A (1.6 oz/5,000 sq ft) of Mainspring Xtra (0.175 lb of thiamethoxam/A and 0.175 lb of cyantraniliprole/A) per year.
- For transplants grown indoors, do not exceed 14.0 oz/A (1.6 oz/5,000 sq ft) of Mainspring Xtra (0.175 lb of thiamethoxam/A and 0.175 lb of cyantraniliprole/A) per crop.
- Do not apply more than 0.2625 lb ai/A thiamethoxam or 0.4 lb ai/A cyantraniliprole per calendar year regardless of formulation or method of application.
- Allow at least 5 days between applications.
- Pre-Harvest Interval: 1 Day
- Water Volume: Use sufficient water volume to insure thorough coverage of foliage. Do not use less than 10 GPA.

Crop	Pests	Rate	Use Restrictions
Cucurbit Vegetables Chayote Chinese waxgourd Citron melon Cucumber Edible gourd Gherkin <i>Momordica</i> species Muskmelon Pumpkin Squash: summer and winter Watermelon	Aphids Cabbage Looper Corn Earworm Cucumber Beetles ¹ Flea Beetles Leafminers Melonworm Pickleworm Rindworm species complex Tobacco Budworm Whiteflies	5.0–7.0 oz/A or 0.6–0.8 oz/5,000 sq ft	Apply before pests reach damaging levels. Apply higher rates within the listed rate range for heavy infestations.

- For transplants grown outdoors, do not exceed 14 oz/A (1.6 oz/5,000 sq ft) of Main-spring Xtra (0.175 lb of thiamethoxam/A and 0.175 lb of cyantraniliprole/A) per year.
- For transplants grown indoors, do not exceed 14 oz/A (1.6 oz/5,000 sq ft) of Main-spring Xtra (0.175 lb of thiamethoxam/A and 0.175 lb of cyantraniliprole/A) per crop.
- Do not apply more than 0.2625 lb ai/A thiamethoxam or 0.4 lb ai/A cyantraniliprole per calendar year regardless of formulation or method of application.
- Allow at least 5 days between applications.
- Pre-Harvest Interval: 1 day
- Water Volume: Use sufficient water volume to ensure thorough coverage of foliage. Do not use less than 10 GPA.
- Pest Control: ¹Suppression

Crop	Pests	Rate	Use Restrictions
Brassica (Cole) Leafy Vegetables Head & Stem Brassica Broccoli, Broccoli Chinese (gai lon) Brussels sprouts Cabbage Cabbage, Chinese (napa) Cabbage, Chinese mustard (gai choy) Cauliflower Cavalo broccolo Kohlrabi Leafy Brassica Greens Broccoli raab (rapini) Cabbage, Chinese (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens	Alfalfa Looper Aphids Beet Armyworm Cabbage Looper Cabbage Webworm Corn Earworm Diamondback Moth Fall Armyworm Flea Beetles Imported Cabbageworm Leafminers (larvae) Sugarbeet Armyworm Thrips (Foliar Feeding) Whiteflies Yellowstriped Armyworm	5.0–7.0 oz/A or 0.6–0.8 oz/5,000 sq ft	Apply before pests reach damaging levels. Apply higher rates within the listed rate range for heavy infestations.

- For transplants grown outdoors, do not exceed 14 oz/A (1.6 oz/5,000 sq ft) of Main-spring Xtra (0.175 lb of thiamethoxam/A and 0.175 lb of cyantraniliprole/A).
- For transplants grown indoors, do not exceed 14 oz/A (1.6 oz/5,000 sq ft) of Main-spring Xtra (0.175 lb of thiamethoxam/A and 0.175 lb of cyantraniliprole/A) per crop.
- Do not apply more than 0.2625 lb ai/A thiamethoxam or 0.4 lb ai/A cyantraniliprole per calendar year regardless of formulation or method of application.
- Allow at least 5 days between applications.
- Pre-Harvest Interval (PHI):
 - 1 day for Head & Stem Brassica
 - 7 days for Leafy Brassica Greens
- Water Volume: Use sufficient water volume to ensure thorough coverage of foliage. Do not use less than 10 GPA.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in a cool, dry place.

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup, procedures and disposal of wastes.

Container Handling (bag)

Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities.

Container Handling (plastic container)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¹/₄ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Mainspring®, the ALLIANCE FRAME, the Syngenta logo, and the PURPOSE ICON are trademarks of a Syngenta Group Company

Viton™ is a trademark of The Chemours Company FC, LLC

©2025 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-866-796-4368.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1424B-L1A 0225