

FIFRA §2(ee) RECOMMENDATION Control of Thrips in Apples

FIFRA Section 2(ee) recommendation for use in Idaho, Washington, Oregon

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Read the “Limitation of Warranty and Liability” in the label booklet for Cormoran® before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This recommendation is made as permitted under FIFRA Section 2(ee) and this bulletin has not been submitted to or approved by U.S. EPA.

Apple

CROPS	PESTS	PRODUCT RATE PER ACRE	REMARKS
Apple, Crabapple, Loquat, Mayhaw and Cultivars, Varieties and/or hybrids of these.	Thrips	20.0-28.0 fl oz	Apply in a minimum finished spray volume of 100 gallons per acre by ground. Do not apply more than two applications against whiteflies or thrips per season.
Pome Fruit (except Pears) Restrictions: <ul style="list-style-type: none"> • Do not apply more than 0.62 lb Novaluron active ingredient containing products per acre per calendar year. • Do not apply more than 0.6 lb Acetamiprid active ingredient containing products per acre per calendar year. • Do not apply more than 95.0 fl oz of formulated product per acre per season. • Repeat applications if needed to maintain control, but do not make applications less than 12 days apart. • Preharvest interval of 14 days. • Restricted entry interval (REI) of 12 hours. 			

Read the “LIMITATION OF WARRANTY AND LIABILITY” in the label booklet for Cormoran before using.