



CROP MIX Z-M-C

A FORMULATION OF 4.5% S, 4.0% Zn, 1.0% Mn, 1.0% Cu

GENERAL INFORMATION

CM CROP MIX Z-M-C is formulated for field, vegetable, row, fruit, tree, vine, ornamental and turf crops. CM CROP MIX Z-M-C may be applied through drip systems, aircraft, or via ground sprayers (high-volume or low-volume). CM CROP MIX Z-M-C may be applied alone or with fertilizers or pesticides. It may be applied either to the plant foliage or to the soil. CM CROP MIX Z-M-C is compatible with most pesticides and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past.

DIRECTIONS

Mixing Sequence

1. Water
2. CM CROP MIX Z-M-C
3. Pesticide

Always add this product to water or nitrogen solution before adding phosphate material.

CONDITIONS OF SALE

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Weather, crop conditions, timing and methods of applications, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MANUFACTURED BY:

CUSTOM AGRICULTURAL FORMULATORS
P.O. BOX 26104, FRESNO, CALIFORNIA 93729

GUARANTEED ANALYSIS

Sulfur (S).....	4.5%
Zinc (Zn).....	4.0%
Copper(Cu).....	1.0%
Manganese (Mn).....	1.0%
Derived from Zinc Sulfate, Copper Sulfate, Manganese Lignosulfonates.	
Wt/Gal: 9.45 Lbs. @ 68°F Gr/ltr: 1133 g @ 20°C	

RATES FOR USE

Foliar Application: Apply 2 to 4 quarts per acre as needed on a 10-14 day schedule. When rates of over 2 quarts are required, split applications of 2 quarts or less are recommended. Avoid foliar applications of CM CROP MIX Z-M-C when plants are under stress due to lack of moisture.

Soil Application:

Maintenance: 1-3 qt/acre
Mild deficiency: 2-6 qt/acre
High deficiency: 1-3 gal/acre

Aerial Application: Apply 1 quart per 5-10 gallons of water per acre. .

Concentrate Application: Apply the suggested rate of this product in enough water for proper coverage.

Dilute Application: Use up to 2 quarts of this product in a minimum of 25 gallons of water per acre.

Drip Irrigation Application: Apply the suggested rate of this product in multiple applications 10-14 days apart throughout the growing season of the crop.

KEEP OUT OF REACH OF CHILDREN

WARNING



PRECAUTIONARY STATEMENTS

WARNING Hazardous to humans and animals. Causes eye and skin irritations. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide labels.

FIRST AID

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Seek medical attention if skin irritation persists. If this product has been swallowed and the person is conscious, immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all governmental procedures for the disposal of pesticide spray solutions. Rinse container; empty container according to approved local practices. Contact county agricultural commissioner for disposal information.

Information regarding the contents and levels of metals in this product is also available by calling 800-355-4457.