



FIFRA 24(c) Special Local Need Label (SLN)

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WISCONSIN

For Control of Diseases on Long-Season Potatoes

BRAVO ULTREX®

For Agricultural or Commercial Use Only

EPA Reg. No. 66222-277

SLN No. WI-180003

Active Ingredient:

Chlorothalonil (tetrachloroisophalonitrile)..... 82.5%

Other Ingredients..... 17.5%

Total: 100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER – PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

This label is valid until December 31, 2022 or until otherwise amended, withdrawn, cancelled, or suspended

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling and federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on this Supplemental label and the EPA registered label for EPA Reg. No. 66222-277.

It is a violation of Federal law to harm an endangered species. Use of this product may be in counties which have endangered and threatened species present; therefore, applicators should evaluate the areas to be treated and take necessary precautions to avoid harming endangered species or their habitat/environment. Growers who are uncertain of the presence of a protected species should contact the Wisconsin Department of Agriculture, Trade and Consumer Protection at 608-224-4547.

Crop	Diseases (Pathogens)	Lb Product/A (lb ai/A)	Application Directions
Potato (Long-season varieties)	Late Blight <i>(Phytophthora infestans)</i> Early Blight <i>(Alternaria solani)</i> Botrytis vine rot <i>(Botrytis cinerea)</i> Black dot <i>(Colletotrichum coccodes)</i>	Low rate: 0.7 (0.6) -then- Higher rates: 0.9 to 1.36 (0.75 to 1.1)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days). Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur: <ul style="list-style-type: none"> • Vines close between rows • Late blight forecasting measures 18 disease severity values (DSV) • The crop reaches 300 P-days Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe. Bravo Ultrex may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo Ultrex label.

Specific Use Restrictions: Do not apply more than 19.35 pounds of Bravo Ultrex (16 lb ai) per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.

Bravo Ultrex is a registered trademark of ADAMA

24(c) Registrant:
 Makhteshim Agan of North America, Inc. d/b/a
 ADAMA
 3120 Highwoods Blvd., Suite 100
 Raleigh, NC 27604

