

SUPPLEMENTAL LABELING

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**

TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS
WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR, EXCEPT THAT UNCERTIFIED
PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

Syngenta Crop Protection, LLC

P. O. Box 18300
Greensboro, North Carolina, 27419-8300
SCP 1652A-S18A 0822

PARAQUAT DICHLORIDE	GROUP	22	HERBICIDE
---------------------	-------	-----------	-----------

**Gramoxone® SL 3.0
HERBICIDE**

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

Active Ingredients:

Paraquat dichloride*	43.9%
Other Ingredients:	56.1%
Total:	100.0%

*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid containing 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.

Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



**DANGER / POISON
PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)
See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

**DIRECTIONS FOR USE
RESTRICTED USE PESTICIDE**

Tobacco; KY, NC, and TN Only

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Tobacco			
Target Weed	Rate pt/A	Application Timing	Use Directions
Refer to Section 7.0 .	1.6 – 2.5	Spring, prior to tobacco transplant	<p>For use in Kentucky, North Carolina and Tennessee Only for burndown of weeds and cover crops prior to transplant.</p> <p>Use the higher rate on dense populations and/or on larger or harder to control weeds.</p> <p>Apply as a broadcast ground application in a minimum of 10 gal/A spray volume.</p> <p>Use the higher rate on dense populations and/or on larger or harder to control weeds. Apply when weeds or cover crop are actively growing and between 1" to 6" in height. Vegetation 6" or taller may not be controlled.</p> <p>Add either: 1) A nonionic surfactant containing 75% or more surface active agent at 0.25% v/v (2 pt per 100 gallons), of the finished spray volume for ground applications, OR 2) A Crop Oil Concentrate (non-phytotoxic) containing 15-20% approved emulsifier, at 1.0 v/v (1 gallon per 100 gallons) of the finished spray volume.</p> <p>See Section 4.4.4 of the Gramoxone SL 3.0 label for adjuvant recommendations.</p>
<p>Tank-Mix Options:</p> <ul style="list-style-type: none"> Gramoxone SL 3.0 may be tank-mixed with other herbicides registered for this use for improved burndown. Refer to Section 4.4.2 for tank-mix precautions in the Gramoxone SL 3.0 label. 			
<p>Resistance Management:</p> <ul style="list-style-type: none"> Refer to Section 3.1. 			
<p>Precaution:</p> <ul style="list-style-type: none"> Weeds that emerge after the application will not be controlled. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Refer to Section 6.1 for additional product use restrictions. Maximum Single Application Rate: 2.5 pt/A REI: 48 hours 			

4. **Minimum Application Interval:** 7 days
5. **Maximum Annual Rate:** 4.0 pt/A
 - a. **DO NOT** exceed 1.5 lb cation/A/year of paraquat-containing products.
6. **DO NOT** make more than 2 application per year.
7. **DO NOT** graze treated areas of feed treated cover crops to livestock.
8. **Pre-Harvest Interval (PHI):** Harvest crop at normal maturity.

Gramoxone® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2022 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina, 27419-8300