

Specimen Label

FLORYLPICOXAMID	GROUP	21	FUNGICIDE
-----------------	-------	----	-----------



Emiox™

Adavelt™ active

FUNGICIDE

™Trademarks of Corteva Agriscience and its affiliated companies

For disease control in established turfgrass on golf courses, including fairways, tee boxes, and greens.

For Use in the contiguous United States and Hawaii only.

Active Ingredient:

Florylpicoxamid:

(2S)-1,1-bis(4-fluorophenyl)propan-2-yl N-[[3-(acetyloxy)-4-methoxypyridin-2-yl]carbonyl]-L-alaninate.....	10.0%
Other Ingredients	90.0%
Total	100.0%

Emiox is formulated as a suspension concentrate (SC) containing 0.83 lb ai/gal of florylpicoxamid.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

EPA Reg. No. 62719-790

Keep Out of Reach of Children

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of Barrier Laminate, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, and Viton ≥ 14 mils.
- Shoes plus socks

User Safety Requirements:

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

FIRST AID

If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
--------------------------------	--

NOTE: Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies, call the poison control center at 800-222-1222. For general information on this product, call 800-992-5994.

Note to Physician: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses: **DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high-water mark.

DO NOT contaminate water when disposing of equipment washwater or rinsate.

Surface Water Advisory: This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of florylpicoxamid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Endangered and Threatened Species Protection Requirements

Before using this product, you must obtain any applicable Endangered Species Protection Bulletins (Bulletins) within six months prior to or on the day of application. To obtain Bulletins, go to Bulletins Live! Two (BLT) at <https://www.epa.gov/pesticides/bulletins>. When using this product, you must follow all directions and restrictions contained in any applicable Bulletin(s) for the area where you are applying the product, including any restrictions on application timing if applicable. It is a violation of Federal law to use this product in a manner inconsistent with its labeling, including this labeling instruction to follow all directions and restrictions contained in any applicable Bulletin(s). For general questions or technical help, call 1-844-447-3813, or email ESPP@epa.gov.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter treated areas without protective clothing until sprays have dried.

Product Information

Emiox™ fungicide may be used for the control of the diseases listed on this label in established turfgrass, on golf courses including fairways, tee boxes, and greens. Optimum disease control is achieved when this product is applied in a regularly scheduled preventative program.

Use Restrictions

- **DO NOT** apply more than 0.472 fl oz/1000 sq ft (20.6 fl oz/acre or 0.133 lb ai/acre) per application.
- **DO NOT** apply more than 1.6 fl oz/1000 sq ft (70 fl oz/acre or 0.45 lb ai/acre) per year.
- **DO NOT** apply more than 3 applications per acre/year at the maximum application rate.
- **DO NOT** apply more than 2 consecutive applications of this product or with other picolinamide fungicides.
- **DO NOT** apply aerially.
- **For Use in the contiguous United States and Hawaii only.**

Use Precautions

- Carefully read, understand, and follow label use rates and restrictions.
- For proper application, determine the size of the area to be treated, the specified label use rate and the gallonage to be applied to the area.
- Use shorter spray intervals between fungicide applications when environmental conditions favor disease development, with susceptible varieties, or disease infection is high.
- Prepare only the amount of spray solution required to treat the measured area.
- Careful calibration of spray equipment is recommended prior to use.

Integrated Pest (Disease) Management

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when disease forecasting models reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

Resistance Management

This product contains florylpicoxamid, a picolinamide Group 21 fungicide. Any fungal population may contain individuals naturally resistant to this product and other picolinamide fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields without rotation with other fungicides from a different cross-resistance Group. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of this product or other picolinamide fungicides within a growing season sequence with fungicides from a different cross-resistance group that control the same pathogens. Avoid consecutive applications of this product or other picolinamide fungicides in a season.
- Use tank mixtures with fungicides from a different cross-resistance group that are equally effective on the target pest when such use is permitted. Use the recommended application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your local company representative. You can also contact your pesticide distributor or university extension specialist to report resistance.

Spray Drift Management

Mandatory Spray Drift Management Mitigations

Groundboom Application

- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- Select nozzle and pressure that deliver medium or coarser spray droplets as indicated in nozzle manufacturer's catalogues and in accordance with American Society of Agricultural & Biological Engineers standards 572.1 and 641 (ASABE S572 and S641).
- During application, the Sustained Wind Speed, as defined by the National Weather Service (standard averaging period of 2 minutes) must register between 3 and 15 miles per hour.
- Wind speed must be measured at the release height or higher, in an area free from obstructions such as trees, buildings, and farm equipment.
- **DO NOT** apply during temperature inversions.

For Ground Boom Application:

- Spray at the appropriate boom height on nozzle selection and nozzle spacing, but **DO NOT** exceed a boom height of 24 inches above target pest or crop canopy. Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions.
- Wind speed and direction must be measured on location using a windsock or anemometer (including systems to measure wind speed or velocity using application equipment).

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if

applications are made improperly or under unfavorable environmental conditions.

Release Height - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce. Automated boom height controllers are recommended with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.

Controlling Droplet Size — Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential is lowest when wind speeds are 15 mph or less. However, many factors, including droplet size, pressure, and equipment type, determine drift potential at any given wind speed. Note: Local terrain can influence wind patterns. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Non-Target Areas

DO NOT apply this pesticide when the product may drift to non-target areas (i.e., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops).

Compatibility

This product is compatible with most commonly used fungicides (except copper-based fungicides), insecticides, growth regulators, micronutrients, spray additives, and adjuvants. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialist prior to use. When an additive and adjuvant is to be used with this product, it is recommended to use a Chemical Producers and Distributors Association certified additive and adjuvant. Select additives and adjuvants that are recommended and/or registered for your specific use pattern and follow their use directions. To determine the physical compatibility of this product with other products, use a jar test. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions and in the correct mixing order as defined below. Invert the jar containing the mixture several times after adding each product and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jells, oily films or layers, or other precipitates, it is not compatible, and the tank mix combination should not be used. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

Spray Additives

Under some conditions, the use of additives or adjuvants may improve the performance of Emiox. However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence turf tolerance and may not match those under which Corteva has conducted testing. Physical incompatibility, reduced disease control, or turf injury may result from mixing Emiox with other products.

A spreading/penetrating type adjuvant such as a nonionic surfactant, crop oil concentrate, or blend may be added at the adjuvant manufacturer's recommended rates.

Mixing Directions

DO NOT prepare more spray mixture than is required for the immediate operation. Thoroughly clean spray equipment before and after using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use. Rinsate may be disposed of on site, according to label use directions, or at an approved waste disposal facility.

Emiox – Alone

- Add 1/2 - 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add this product to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after the product has completely and uniformly dispersed into the water.
- Maintain agitation until all the mixture has been sprayed.

Emiox – Tank Mix

Mixing Order for Tank Mixes:

1. Fill the spray tank to at least 1/4 to 1/3 of the required spray volume.
2. Start agitation.
3. Add different formulation types in the following order, allowing time for complete mixing and dispersion after addition of each while following label directions for each material added to the tank.
 - (a) Water soluble bags
 - (b) Dry flowables
 - (c) Wettable powders
 - (d) Suspensions and liquid flowables
 - (e) This product and other emulsifiable concentrates and any solutions
 - (f) Adjuvants
4. Finish filling the spray tank. Maintain continuous agitation during mixing, final filling and throughout application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

If application or agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be re-suspended before spraying is resumed.

Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix product label. Label dosage rate must not be exceeded, and the most restrictive label

precautions and limitations must be followed. This product must not be mixed with any product which prohibits such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Recommended Clean-out Procedures for Spray Equipment:

1. Drain any remaining spray mixture from the application equipment, then wash out tank, boom, and hoses with clean water. Drain again.
2. Hose down the interior surfaces of the tank while filling the tank 1/2 full of water.
3. Add commercial tank cleaner at manufacturers recommended rates, or at a rate of 1 gallon per 100 gallons of water. Recirculate for 10 - 20 minutes and spray out the mixture through the boom.
4. Remove all spray nozzles and screens and clean separately.
5. Thoroughly clean exterior surfaces of spray equipment.

Application Instructions

Apply this product in conjunction with management practices for crop uses that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Use diagnostic kits, extension experts, accredited laboratories, or other identification methods when developing disease control strategies.

For optimum disease control, apply this product preventatively in effective disease management programs following the specified label rates, use directions, use restrictions, and resistance management strategies. For best results, allow this product to dry at least 1 hour before any rainfall or field irrigation.

Ground Application

Apply this product in sufficient water volume to ensure thorough coverage of the canopy for optimum disease control. Refer to Specific Uses and Directions for additional information.

Air Blast Application

Air-assisted or air-blast sprayers move spray droplets into the crop canopy using a forced-air system. The fan must be set up to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area. Equip sprayers with nozzles that provide accurate and uniform application.

USES

Established Turfgrass on Golf Courses

Disease Controlled or Suppressed	Rate (fl oz/1000 sq ft)	Rate (fl oz/acre)	Rate (lb ai/acre)	Application Interval (Days)	Application Instructions
Dollar Spot <i>Clarireedia homoeocarpa</i> , <i>Clarireedia</i> spp. (formerly known as <i>Sclerotinia homoeocarpa</i>) Anthracnose (<i>Colletotrichum graminicola</i>)	0.317-0.472	13.8-20.5	0.09-0.133	14 to 21*	Apply preventatively when conditions are favorable for disease development

Restrictions:

- **DO NOT** apply more than 0.472 fl oz/1000 sq ft (20.5 fl oz/acre or 0.133 lb ai/acre) per application.
- **DO NOT** apply more than a total of 1.6 fl oz/1000 sq ft per year (70 fl oz/acre or 0.45 lb ai/acre)
- **DO NOT** apply more than 5 applications per acre/year at the minimum application rate, 3 applications per acre/year at the maximum application rate, or 1/3 of total sprays in a program, whichever is the lower.
- **DO NOT** tank mix with copper-based fungicides.
- **DO NOT** apply aerially.
- For golf courses, use only on tees, greens, and fairways. **DO NOT** use florylpicoxamid containing products on roughs.
- **Minimum Retreatment Interval:** 14 days
- **DO NOT** apply more than 2 consecutive applications of this product or with other picolinamide fungicides.
- Not for homeowner use.
- For use in the contiguous US and Hawaii only.

*Under favorable conditions for high disease pressure use the shorter application interval.

(Storage and Disposal for rigid containers 5 gal or less)

Storage and Disposal

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

(Storage and Disposal for refillable rigid containers greater than 5 gal)

Storage and Disposal

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

(Storage and Disposal for non-refillable rigid containers larger than 5 gal)

Storage and Disposal

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation, of Remedies.

Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions for use, subject to the inherent risks set forth below. To the extent consistent with applicable law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. Corteva Agriscience will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by Corteva Agriscience. To the extent consistent with applicable law, all such risks associated with non-directed use shall be assumed by buyer and/or user.

Limitation of Remedies

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent consistent with applicable law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent consistent with applicable law, in no case shall Corteva Agriscience be liable for consequential, incidental or special damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

TM® Trademarks of Corteva Agriscience and its affiliated companies

Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268

Label Code: CD02-332-020
Initial Publication

EPA accepted 7/10/2026

Manténgase fuera del alcance de los niños

CAUTION PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DECLARACIONES PREVENTIVAS

Riesgos para los seres humanos y animales domésticos

PRECAUCIÓN

Dañino si se absorbe a través de la piel. Evite el contacto con la piel, los ojos o la ropa. Lávese con abundante agua y jabón luego de manejar (pesticidas) y antes de comer, beber, masticar chicle o tabaco, o ir al baño. Quítese y lave la ropa contaminada antes de reutilizarla.

Equipo de protección personal (PPE)

Los aplicadores y otros manipuladores de pesticidas deberán usar:

- Camisa de manga larga y pantalones largos.
- Guantes resistentes a productos químicos fabricados con laminados impermeables, goma de nitrilo ≥ 14 milésimas de pulgada, goma de neopreno ≥ 14 milésimas de pulgada y Viton ≥ 14 milésimas de pulgada.
- Zapatos y calcetines.

Requisitos de seguridad para el usuario:

Siga las instrucciones del fabricante para la limpieza/mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE separadamente de otra ropa para lavar.

PRIMEROS AUXILIOS

Si cae en la piel o la ropa:

- quítese la ropa contaminada.
- Enjuague la piel inmediatamente con abundante agua durante 15-20 minutos.
- Llame a un centro de control de envenenamientos o a un médico para consejo de tratamiento.

NOTA: cuando llame a un centro de control de envenenamientos, o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. En caso de emergencia médica, llame al centro de control de envenenamientos al 800-222-1222. Para obtener información general sobre este producto, llame al 800-992-5994.

Nota al médico: contiene destilado de petróleo. Los vómitos pueden provocar neumonía por aspiración.

Recomendaciones de seguridad para el usuario

Los usuarios deben:

- Quitarse de inmediato la ropa/el PPE si entra pesticida en su interior. Luego, deben lavarse por completo y ponerse ropa limpia.
- Quitarse de inmediato el PPE después de manipular este producto. Lavar la parte externa de los guantes antes de quitárselos. Tan pronto como sea posible, lavarse con abundante agua y ponerse ropa limpia.

Riesgos ambientales

Para el uso terrestre: **NO** aplique directamente en el agua, en zonas donde haya aguas superficiales ni en zonas intermareales por debajo de la marca de agua máxima promedio. **NO** contamine el agua cuando deseche aguas de lavado o de enjuague del equipo.

Recomendación para aguas superficiales: Este producto puede afectar la calidad de las aguas superficiales debido a la escorrentía del agua de lluvia. Esto es especialmente cierto en tierra con drenaje escaso y con aguas subterráneas poco profundas.

Una franja de protección vegetal nivelada y bien conservada entre las áreas en las que se aplica este producto y las fuentes de aguas superficiales, como estanques, arroyos y manantiales, reducirá la carga potencial de florilpicoxamida del agua de escorrentía y los sedimentos. La escorrentía de este producto se reducirá si se evitan las aplicaciones cuando se espera que llueva o se riegue en un plazo de 48 horas.

INSTRUCCIONES DE USO

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal. Lea atentamente todas las Instrucciones de uso antes de aplicarlo.

NO aplique este producto de forma que entre en contacto con trabajadores u otras personas, ya sea directamente o a través de la deriva. Solo los manipuladores de pesticidas con la protección adecuada pueden permanecer en el área durante la aplicación. Para obtener información sobre cualquier requisito específico de su estado o tribu, consulte a la agencia responsable de la regulación de pesticidas.

Requisitos de protección de especies en peligro de extinción y amenazadas

Antes de utilizar este producto, debe obtener el Boletín de Protección de Especies en Peligro de Extinción (Boletín) correspondiente dentro de los seis meses anteriores o el mismo día de la aplicación. Para obtener los boletines, visite Bulletins Live! Two (BLT) en <https://www.epa.gov/pesticides/bulletins>. Al utilizar este producto, debe seguir todas las instrucciones y restricciones que figuren en el boletín correspondiente a la zona donde aplique el producto, incluidas las restricciones sobre el momento de la aplicación, si las hubiera. El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal, incluida esta instrucción de la etiqueta de seguir todas las indicaciones y restricciones incluidas en el Boletín correspondiente. Si tiene preguntas generales o necesita asistencia técnica, llame al 1-844-447-3813 o envíe un correo electrónico a ESPP@epa.gov.

Requisitos para usos no agrícolas

Los requisitos en esta sección de la etiqueta aplican a los usos de este producto que **NO** están cubiertos por el Estándar para la Protección del Trabajador Agrícola para pesticidas agrícolas (40 CFR, Parte 170). El WPS se aplica cuando este producto se utiliza para producir plantas agrícolas en granjas, bosques, viveros o invernaderos.

NO ingrese a las áreas tratadas sin vestimenta de protección hasta que las pulverizaciones se hayan secado.

(Almacenamiento y desecho de envases rígidos con una capacidad igual o inferior a 5 galones)

Almacenamiento y desecho

NO contamine el agua, la comida ni los forrajes mediante el almacenamiento y desecho.

Almacenamiento del pesticida: almacene en el envase original.

En caso de fuga o derrame, contenga el material y deséchelo como residuo.

Desecho de pesticidas: los residuos resultantes del uso de este producto deben desecharse en el lugar o en un centro de desecho de residuos autorizado.

Manipulación del envase: envase no rellenable. **NO** reutilice ni rellene este envase. Ofrezcalo para ser reciclado, si esa opción está disponible, o perforélo y deséchelo en un relleno sanitario, o mediante otros procedimientos permitidos por las autoridades estatales y locales. Enjuague el envase (o equivalente) tres veces o a presión inmediatamente después de vaciarlo. **Enjuague tres veces** de la siguiente manera: vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla y drene durante 10 segundos después de que el flujo comience a gotear. Llene el envase hasta 1/4 de su capacidad con agua y vuelva a taparlo. Agite durante 10 segundos. Vierta el agua de enjuague en el equipo de aplicación o en un tanque de mezcla o almacene las aguas de enjuague para su uso o desecho posterior. Drene durante 10 segundos después de que el flujo comience a gotear. Repita este procedimiento dos veces más. **Enjuague a presión** de la siguiente manera: vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla y continúe drenando durante 10 segundos después de que el flujo comience a gotear. Mantenga el envase boca abajo sobre el equipo de aplicación o el tanque de mezcla o recoger las aguas de enjuague para su uso o desecho posterior. Inserte la boquilla de enjuague a presión en el lateral del envase y enjuague a unos 40 psi durante al menos 30 segundos. Drene durante 10 segundos después de que el flujo comience a gotear. Luego, ofrezca el envase para ser reciclado, si esa opción está disponible, o perforélo y deséchelo en un relleno sanitario, o mediante otros métodos aprobados por las autoridades estatales y locales.

(Almacenamiento y desecho de contenedores rígidos rellenos de más de 5 galones)

Almacenamiento y desecho

NO contamine el agua, la comida ni los forrajes mediante el almacenamiento y desecho.

Almacenamiento del pesticida: almacene en el envase original.

En caso de fuga o derrame, contenga el material y deséchelo como residuo.

Desecho de pesticidas: los residuos resultantes del uso de este producto deben desecharse en el lugar o en un centro de desecho de residuos autorizado.

Manipulación del envase: envase rellenable. Rellene este envase solo con pesticidas. **NO** reutilice este envase para ningún otro fin.

La limpieza del envase antes de su disposición final es responsabilidad de la persona que deseché el envase. La limpieza antes de rellenarlo es responsabilidad de la persona que lo rellena. Para limpiar el envase antes de su disposición final, vacíe el contenido restante de este envase en el equipo de aplicación o en el tanque de mezcla. Llene el envase con un 10 % de agua. Agite enérgicamente o haga recircular el agua con la bomba durante dos minutos. Vierta o bombee el agua de enjuague en el equipo de aplicación o en el sistema de recolección de aguas de enjuague. Repita este procedimiento de enjuague dos veces más. Luego, ofrezca el envase para ser reciclado, si esa opción está disponible, o perfórelo y deséchelo en un relleno sanitario, o mediante otros métodos aprobados por las autoridades estatales y locales.

(Almacenamiento y desecho de contenedores rígidos no rellenos con una capacidad superior a 5 galones)

Almacenamiento y desecho

NO contamine el agua, la comida ni los forrajes mediante el almacenamiento y desecho.

Almacenamiento del pesticida: almacene en el envase original.

En caso de fuga o derrame, contenga el material y deséchelo como residuo.

Desecho de pesticidas: los residuos resultantes del uso de este producto deben desecharse en el lugar o en un centro de desecho de residuos autorizado.

Manipulación del envase: envase no rellenable. **NO** reutilice ni rellene este envase. Ofrézcalo para ser reciclado, si esa opción está disponible, o perfórelo y deséchelo en un relleno sanitario, o mediante otros procedimientos permitidos por las autoridades estatales y locales.

Enjuague el envase (o equivalente) tres veces o a presión inmediatamente después de vaciarlo. **Enjuague tres veces** de la siguiente manera: vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla. Llene el envase hasta 1/4 de su capacidad con agua. Vuelva a colocar los cierres y ajústelos. Coloque el envase sobre un lado, hágalo rodar hacia adelante y hacia atrás, y asegúrese de que dé al menos una vuelta completa, durante 30 segundos.

Coloque el envase sobre su extremo e inclínelo hacia adelante y hacia atrás varias veces. Dé vuelta el envase, colóquelo sobre su otro extremo e inclínelo hacia adelante y hacia atrás varias veces. Vacíe las aguas de enjuague en el equipo de aplicación o en un tanque de mezcla, o bien, almacene las aguas de enjuague para su uso o desecho posterior.

Repita este procedimiento dos veces más. **Enjuague a presión** de la siguiente manera: vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla y continuar drenando durante 10 segundos después de que el flujo comience a gotear. Mantenga el envase boca abajo sobre el equipo de aplicación o el tanque de mezcla o recoger las aguas de enjuague para su uso o desecho posterior. Inserte la boquilla de enjuague a presión en el lateral del envase y enjuague a unos 40 psi durante al menos 30 segundos. Drene durante 10 segundos después de que el flujo comience a gotear. Luego, ofrezca el envase para ser reciclado, si esa opción está disponible, o perfórelo y deséchelo en un relleno sanitario, o mediante otros métodos aprobados por las autoridades estatales y locales.