



D & M Chem

7 PAC LIQUID

GUARANTEED ANALYSIS

Magnesium (Mg) Total.....	1.5%
Boron (B)38%
Copper (Cu)13%
Iron (Fe)	1.5%
Manganese (Mn)	1.5%
Molybdenum (Mo)08%
Zinc (Zn)	1.13%

Derived from: Copper, Iron, Magnesium, Manganese and Zinc are chelated by Citric acid and EDTA Chelate (Disodium Ethylenediamine Tetra Acetate Chelate). Boron derived from Sodium Borate and Molybdenum derived from Sodium Molybdate.

Information regarding the contents and levels of metals in this product is available on the Internet at www.aapfco.org/metals.htm.

KEEP OUT OF REACH OF CHILDREN

DANGER

GHS Hazards: Harmful if swallowed. Causes severe skin burns and eye damage. May cause damage to organs through prolonged or repeated exposure. **GHS Precautions:** Wash face, hands and any exposed skin thoroughly after handling.

Do not eat, drink or smoke when using this product. Do not breathe dust/fume/gas/mist/vapors/spray. Wear protective gloves/protective clothing/eye protection/face protection.

FIRST AID

Immediately call a poison center or doctor/physician.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a poison center or doctor/physician.

IF ON SKIN (or hair): Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower. Wash contaminated clothing before reuse.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Immediately call a poison center or doctor/physician.

IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell. Rinse mouth. Do not induce vomiting.

NET WEIGHT: 10.8 POUNDS PER/GALLON

NET CONTENTS: 2.5 Gallons or Bulk



D & M Chem

7 PAC LIQUID

FOR TREATMENT OF SOILS AND CROPS WHERE MICRONUTRIENT DEFICIENCY IS DIAGNOSED OR SUSPECTED.

7 PAC LIQUID is particularly suitable for foliar application, inclusion in liquid feed and hydroponic mixes and for soil application. Foliar fertilization is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by agricultural crops. Plants need only very small amount of micronutrients for maximum growth. Thus, while a deficiency of any essential element will greatly reduce plant growth, the overuse of micronutrients can produce a harmful level of these nutrients in the soil which may be more difficult to correct than deficiency. This fertilizer is to be used only on crops which respond to Molybdenum. Crops high in Molybdenum are toxic to grazing animals.

Boron: This fertilizer contains Boron, which may be injurious to certain crops. Contact your local county agent or field consultant for specific information.

APPLICATION RATES

Use 1 qt to 2 qts For crops such as alfalfa, almonds, avocados, apples, beans (all), broccoli, cabbage, cauliflower, carrots, celery, citrus, corn, grapes, hops, lettuce, milo, melons, nectarines, rice, pears, peaches, pecans, peppers, plums, prunes, potatoes, peanuts, sorghum, sugar beets, sweet corn, sugar cane, strawberries, tomatoes, turnips, walnuts, watermelons, wheat and most other crops.

ORNAMENTALS: Two teaspoons per gallon of water every 2 to 4 weeks. Use enough water to cover thoroughly or approximately 1 gallon per 10 square feet. 7 PAC LIQUID can be incorporated into NPK fertilizer at 8 oz. per 50 lbs. of NPK.

APPLICATION RECOMMENDATION FOR TURF:

GREENS: Apply 7 PAC LIQUID 1 qt per acre (43,650 sq. ft.) in a minimum of 150 to 200 gallons of water. 7 PAC LIQUID at 1 qt. in 150 to 200 gallons of water will cover approximately six average green. Best results will be seen where 7 PAC LIQUID is applied in early spring when green are actively growing. Four or more applications may be made at monthly intervals to help maintain vigor and color. Applications should always be made later in the evening or early in the morning to reduce potential for phytotoxicity.

FAIRWAYS AND LAWNS: Apply 7 PAC LIQUID at 1 qt per acre in a minimum of 40 gallons of water. Use three or more applications per year to maintain fine healthy turf.

CAUTION: Do not apply more than 2 qts of any single chelate or mixed chelates per foliar application.

NOTICE

Manufacturer warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Manufacturer. In no case shall Manufacturer be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Manufacturer makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

FOR INFORMATION CONTACT

D & M CHEM INC.
112 CHARRON ROAD
MOXEE, WA 98936
(509) 454-8154