

# SUPPLEMENTAL LABEL



PYRAFLUFEN-ETHYL	GROUP	14	HERBICIDE
------------------	-------	----	-----------

EPA Reg. No. 71711-25

Crop: Hops

This supplemental label expires September 30, 2022 and must not be used or distributed after this date.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application.

Read the label affixed to the container for VENUE® Herbicide before applying. Use of VENUE Herbicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for VENUE Herbicide. New use directions appear on this supplemental label that may be different from those that appear on the container label.

## CROP USE DIRECTIONS

Hops			
Application	Pest	Rate/Acre	Application Instructions
In-Season	Listed Broadleaf Weeds  Sucker Management	3.0 to 4.0 fl oz/acre  (0.0040 to 0.0053 lb ai/acre)	<ul style="list-style-type: none"> <li>Apply in a minimum of 20 gallons spray solution per acre by ground equipment to target weeds and sucker growth.</li> <li>Use of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions.</li> <li>COC adjuvants are advised, although other adjuvants may be used.</li> <li>Use the higher rate for hard-to-control weeds.</li> <li>For the management of undesirable sucker growth on the basal portion of trunks, root sprouts, and vine trunks, growth must be controlled when the tissue is young, immature, and/or not hardened off.</li> <li>Avoid contact with green, uncallused bark of young trees or vines established less than one year unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers.</li> <li>Direct application to the lower 12-18 inches of the hops plant and extend out to approximately 20-40 inches from each side of the row.</li> <li>Calculate the rate on a treated acre basis as follows: Assuming a plot length of 50 feet: Calculations would be 1.5 feet (if 18-inch extension), 2 feet (if 24-inch extension) x 50 feet x 2 passes (one on each side of the vine).</li> </ul> <p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> <li><b>Do not</b> apply by air for this use.</li> <li><b>Do not</b> allow spray to drift onto desirable fruit, foliage, vines, or trees as damage will occur.</li> <li><b>Do not</b> exceed 4.0 fl oz (0.0053 lb ai) per acre in a single application for this use.</li> <li><b>Do not</b> exceed a combined total of 2 applications per year for this use.</li> <li>Allow a minimum of 30 days between applications for this use.</li> <li><b>Do not</b> exceed 8.0 fl oz (0.0106 lb ai) per acre per year for all applications combined.</li> <li>Preharvest Interval (PHI): 30 days</li> </ul>

©2019 Nichino America, Inc. Venue and Nichino America logo are registered trademarks of Nichino America, Inc.

**NICHINO**  
**AMERICA**

Nichino America, Inc.  
4550 Linden Hill Road, Suite 501  
Wilmington, DE 19808  
888-740-7700