

SUPPLEMENTAL LABELING

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**
TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS
WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR, EXCEPT THAT UNCERTIFIED
PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina, 27419-8300
SCP 1652A-S16A 0822

PARAQUAT DICHLORIDE	GROUP	22	HERBICIDE
---------------------	-------	-----------	-----------

Gramoxone® SL 3.0
HERBICIDE

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

Active Ingredients:

Paraquat dichloride*	43.9%
Other Ingredients:	56.1%
Total:	100.0%

*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid containing 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.

Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



DANGER / POISON
PELIGRO / VENENO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)
See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



**DIRECTIONS FOR USE
RESTRICTED USE PESTICIDE**

Strawberries, FL Only

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Strawberries			
Target Weed	Rate pt/A	Application Timing	Use Directions
Refer to Section 7.0.	1.3	Post-Harvest	<p>For use in Florida Only for desiccation of strawberry plants following harvest.</p> <p>Apply as a ground broadcast application in a minimum of 30 gal/A spray volume.</p> <p>Always use one of the following: 1) A nonionic surfactant containing 75% or more surface-active agent at 0.125% v/v (1 pint per 100 gallons) or a nonionic surfactant containing 50-74% surface-active agent at 0.25% v/v (2 pints per 100 gallons), of the finished spray volume, OR 2) A crop oil concentrate (nonphytotoxic) containing 15-20% approved emulsifier, at 1.0% v/v (1 gal/100 gallons) of the finished spray volume.</p> <p>Use flat fan nozzles for the most effective application.</p> <p>See Section 4.4.4 of the Gramoxone SL 3.0 label for adjuvant recommendations.</p>
Tank Mix Options:			
<ul style="list-style-type: none"> • Gramoxone SL 3.0 can be tank mixed with other herbicides labeled for use on strawberries applied by the same methods and at the same timings. Refer to Section 4.4.2 for tank-mix precautions in the Gramoxone SL 3.0 label. 			
Resistance Management:			
<ul style="list-style-type: none"> • Refer to Section 3.1. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1. Refer to Section 6.1 for additional product use restrictions. 2. Maximum Single Application Rate: 1.3 pt/A 3. REI: 48 hours 4. Minimum Application Interval: 7 days 5. Maximum Annual Rate: 3.9 pt/A <ol style="list-style-type: none"> a. DO NOT exceed 1.5 lb cation/A/year of paraquat-containing products. 6. DO NOT make more than 3 application per year. 7. DO NOT use treated fruit for human or animal consumption. 8. DO NOT apply when weather conditions favor drift from treated areas. 9. DO NOT apply this product through any type of irrigation system. 10. DO NOT graze livestock in treated area. 11. Pre-Harvest Interval (PHI): For use only on fields where harvest operations have finished for season. 			

Gramoxone® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2022 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina, 27419-8300