

PILATUS® FOL

1-0-5

GUARANTEED ANALYSIS

Total Nitrogen (N)	1.0%
1.0% Water-Soluble Nitrogen	
Soluble Potash (K ₂ O)	5.0%
Boron (B)	0.09%
Manganese (Mn)	0.18%
Zinc (Zn)	0.41%

Derived from: corn protein hydrolysate, muriate of potash, disodium octaborate, manganese sulfate, and zinc sulfate monohydrate.

WARNING: SOME CROPS MAY BE INJURED BY APPLICATION OF BORON

KEEP OUT OF REACH OF CHILDREN

For Product Use Information: 1-866-761-9397
SDS is available at www.arysta-na.com

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.aapfco.org/metals.htm>

F1721

Produced for:

Arysta LifeScience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

NET WEIGHT: 23 Pounds (10.42 kg)
NET CONTENTS: 2.5 Gallons (9.46 liters)

AD110415V7

Application instructions: Apply **PILATUS® FOL** as a foliar spray in enough water to thoroughly cover the foliage, usually 10 – 100 gallons per acre for ground equipment and 5 -10 gallons per acre for aerial application. Refer to the Table below for application rates and timing.

CROP	RATE/ACRE	TIMING
Banana	16-24 fl oz	Apply 30-45 days after transplant. Repeat monthly.
Cereal Grains, such as: Barley, Corn, Millet, Oats, Popcorn, Rice, Rye, Sorghum, Wheat, Wild Rice	16 fl oz	Apply in small grains at the flag leaf stage (Feekes 9 to 10). Apply to Corn and Sorghum at the V-4 to V-7 stage.
Citrus, such as: Calamondin, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin, Orange (sour and sweet), Pummelo, Tangerine	16 fl oz/ 100 gal water	Apply two weeks after bloom begins. Repeat during bloom and at petal drop.
Cotton	16 fl oz	Apply at early bloom. Repeat 2 to 3 weeks after first spray.
Cole Crops, such as: Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage (Chinese), Cauliflower, Cavalo Braccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens	16-24 fl oz	Apply when there are 2 to 3 pairs of leaves. Repeat 30-45 days after first spray.
Cucurbit Vegetables, such as: Cantaloupe, Chayote, Chinese Waxgourd, Citron Melon, Cucumbers, Gherkin, Edible Gourds, Muskmelon hybrids and/or cultivars, Pumpkin, Summer Squash (such as Crookneck and Zucchini), Watermelons, Winter Squash (such as Acorn and Butternut)	16 fl oz	Apply at the beginning of bloom and Repeat 2 times at 14 day intervals.
Fruiting Vegetables, such as: Eggplant, Groundcherry, Pepino, Peppers (such as Bell, Chili, Pimento and Sweet Peppers), Tomatillo, and Tomatoes	16 fl oz	Apply at the beginning of bloom. Repeat 2 times at 14 day intervals.
Grapes, such as: Dried, Table, Wine	24-28 fl oz	For seedless grapes, apply after blooming. Repeat 3-4 times at weekly intervals.
	24- 32 fl oz	For Superior and Red Globe varieties, apply to bunches. Repeat 1 to 2 times at weekly interval.
Leafy Vegetables, such as: Amaranth, Arugula, Asparagus, Cardoon, Celery, Celtuce, Chervil, Chrysanthemum, Corn Salad, Cress (garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio, Rhubarb, Spinach, Swiss Chard.	16-20 fl oz	Apply when there are two to three pair of leaves and repeat 10 – 15 days later.
Pineapples	24-30 fl oz/ 250 gal water	Apply 18 days after flora induction. Repeat 24 days later.
Pome Fruit, such as: Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental), Quince		Apply between bloom and petal fall. Repeat when fruit appears.

CROP	RATE/ACRE	TIMING
Potatoes, Tuber Vegetables, such as: Arracacha, Arrowroot, Artichoke, Canna (edible), Cassava, Chayote (root), Garlic, Ginger, Onion, Potato, Sweet Potato, Tanier, Turmeric, Yam Bean, Yam	16-24 fl oz	Apply during bloom. Repeat 2 times at 14 day intervals.
Root Vegetables, such as: Beet (garden & sugar), Burdock (edible), Carrot, Celeriac, Chervil, Chicory, Ginseng, Horseradish, Parsnip, Radish, Rutabaga, Salsify, Skirret, Turnip	16-24 fl oz	Apply during bloom. Repeat 15-20 days later.
Strawberries	16 fl oz	Apply when blooming begins. Repeat every 21 day for a total of 8 applications.
Stone Fruit, such as: Apricot, Cherry (sweet and tart), Nectarine, Peach, Plum, Plumcot, Prune (fresh)	12-16 fl oz/ 100 gal water	Apply between bloom and petal fall. Repeat when fruit appears.
Succulent, Dried Beans and Peas, such as: Beans (<i>Phaseolus</i> and <i>Vigna</i> spp.), Jackbean, Pea (<i>Pisum</i> spp.), Pigeon Pea, Soybean, Sword Bean, and dried cultivars of Bean (<i>Lupinus</i> , <i>Phaseolus</i> , <i>Vigna</i> spp.), Broad Bean, Chickpea, Guar, Lablab Bean, Lentil, and Pea.	16 fl oz	Apply one application between R1 and R3 growth stage.
Tree Nuts and Pistachio, such as: Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory Nut, Macadamia Nut, Pecan, Walnut (Black and English), and Pistachio	12-16 fl oz/ 100 gal water	Apply between bloom and petal fall. Repeat when fruit appears.
Tropical Tree Fruits, such as: Avocado, Cherimoya, Coffee, Guava, Lychee, Mango, Papaya	12-16 fl oz/ 100 gal water	Apply 2 weeks after bloom begins. Repeat during bloom and at petal drop.

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Arysta LifeScience North America, LLC ("Arysta"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

Arysta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Arysta, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA, MANUFACTURER, OR THE SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE BUYER AND THE USER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF ARYSTA IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER OR USER'S EXCLUSIVE REMEDY, AND ARYSTA'S, THE MANUFACTURER'S, OR THE SELLER'S TOTAL LIABILITY SHALL BE LIMITED TO DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT, OR AT ARYSTA'S ELECTION, THE REPLACEMENT OF THE PRODUCT.

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.

PILATUS is a registered trademark of Arysta LifeScience North America, LLC.