



PROSPER[®]

EverGol[®]

For use as a fungicide and insecticide seed dressing for protection against listed soilborne, seedborne, and post-emergence diseases and insect pests of listed crop plants.

ACTIVE INGREDIENTS:

Clothianidin	22.30%
Penflufen	0.82%
Trifloxystrobin	0.55%
Metalaxyl	0.55%

OTHER INGREDIENTS: 75.78%

Contains: 2.42 lb clothianidin; 0.09 lb penflufen; **TOTAL 100.00%**

0.06 lb trifloxystrobin; 0.06 lb metalaxyl per gallon

Contains: 290 g clothianidin; 10.7 g penflufen; 7.15 g trifloxystrobin; 7.15 g

metalaxyl per liter

EPA Reg. No. 264-1121

CLOTHIANIDIN	GROUP 4A	INSECTICIDE
PENFLUFEN	GROUP 7	FUNGICIDE
TRIFLOXYSTROBIN	GROUP 11	FUNGICIDE
METALAXYL	GROUP 4	FUNGICIDE

KEEP OUT OF REACH OF CHILDREN CAUTION

For **MEDICAL** and **TRANSPORTATION** Emergencies

ONLY Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER

(1-866-992-2937)

**Please refer to booklet for additional precautionary
statements and directions for use.**

Produced for:
Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167

PROSPER and EVERGOL are registered trademarks of Bayer.
©2019 Bayer CropScience

190923D 11/19

US80523157D

FIRST AID

IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) or viton ≥ 14 mils.

All mixers, loaders, applicators and other handlers must wear:

- long sleeved shirt and long pants,
- socks plus shoes,
- chemical resistant gloves, except when bagging or sewing bags of treated seeds.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to bees exposed directly (contact) and is toxic to birds, mammals, fish and aquatic invertebrates. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or run-off from treated areas is hazardous to fish and aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seed spilled during loading.

This product has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use only in commercial seed treatment equipment. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.

PRODUCT INFORMATION

Prosper® EverGol®:

- is an insecticide and fungicide seed treatment for use on canola (including *Brassica napus* and *Brassica rapa*), mustard (oilseed and condiment) and rapeseed for control of listed insect pests and for control of seed rot, damping off, seedling blight and early season root rot caused by *Pythium*, *Rhizoctonia*, *Fusarium* and *Alternaria*.
- also controls seedborne Blackleg.
- insecticide component, clothianidin, controls flea beetles on seeds and seedlings of canola (including *Brassica napus* and *Brassica rapa*), mustard (oilseed and condiment) and rapeseed. For protection of seeds and emerging seedlings only.

RESTRICTIONS

- Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: Do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

RESISTANCE MANAGEMENT RECOMMENDATION

For resistance management, please note that Prosper EverGol contains a Group 4 [Phenylamides class], a Group 7 [Succinate-dehydrogenase inhibitor class] and a Group 11 [Quinone outside inhibitor class] fungicide. Any fungal population may contain individuals naturally resistant to Prosper EverGol and other Group 4, 7, or 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Prosper EverGol contains a Group 4A [Neonicotinoids class] insecticide. Any insect population may contain individuals naturally resistant to Prosper EverGol and other Group 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide and insecticide resistance, take one or more of the following steps:

- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

COMMERCIAL SEED TREATMENT

CROP	DISEASE/PEST	APPLICATION RATE	DIRECTIONS
Canola (including <i>Brassica napus</i> and <i>Brassica rapa</i>), Rapeseed and Mustard (oilseed and condiment)	For control of listed insect pests and for control of seed rot, damping off, seedling blight and early season root rot caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> and <i>Alternaria</i> . Also controls seedborne Blackleg. The insecticide component, clothianidin, controls flea beetles on seeds and seedlings of canola (including <i>Brassica napus</i> and <i>Brassica rapa</i>), mustard (oilseed and condiment) and rapeseed. For protection of seeds and emerging seedlings only.	Apply 21.5 fl oz of Prosper EverGol to 100 lbs of seed (1.4 liters of product per 100 kgs of seed) (200.8 g ai per 100 lb seed, based on an average of 100,000 seeds per lb) Maximum number of application per year is 3.	Apply using commercial slurry or mist-type seed treatment equipment. Mix thoroughly before use. Use high quality, properly cleaned seed. Pre-test all tank mixes to determine the physical compatibility between formulations. Follow all cautions and most restrictive limitations on labeling for all products used in mixtures. Prior to and during application, Prosper EverGol must be thoroughly agitated to ensure uniform mixing.

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with Prosper EverGol must be labeled with the following statements:

- This seed has been treated with Prosper EverGol which contains metalaxyl, clothianidin, trifloxystrobin, and penflufen.
- Do not use treated seed for food, feed, or oil production. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- User is responsible for ensuring that the seed bag meets all the requirements under the Federal Seed Act.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with Prosper EverGol:

- Store away from feeds and foodstuffs.
- Wear long sleeved shirt, long pants, shoes, socks, and chemical resistant gloves when handling treated seed.

(continued)

- Areas planted with treated seed may be replanted immediately with canola, cotton, corn (field, pop and sweet), cereal grains except rice, mustard, rapeseed, sorghum, soybean, and potatoes (tuberous and corm subgroup 1C). These areas may also be replanted no less than 30 days after planting treated seeds with grasses, non-grass livestock feeds, dried beans, fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables) and sugar beets. Do not plant any other crop in the treated area for at least twelve months after treated seeds are planted.
- After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows worker to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or planting media.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- This contains metalaxyl that is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.
- Dispose of seed packaging in accordance with local requirements.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch.
- This compound is toxic to birds and mammals. Treated seed exposed on soil surface may be hazardous to birds and mammals. Cover or collect seeds spilled during loading.
- This compound may be highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- Regardless of type of application, do not apply more than 0.2 lbs of the active ingredient clothianidin per acre per season.
- For Canola (including *Brassica napus* and *Brassica rapa*), rapeseed and Mustard (based on canola, rapeseed or mustard seeding rate): Do not exceed 0.00123 lb. penflufen/A (0.6 g ai/A) per crop season. The maximum number of applications per year is 3.
- Runoff may be hazardous to aquatic organisms in water adjacent to treated areas.
- This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Store in original container away from feed and food. Store in cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 105°F (40°C) or in temperatures that fall below 14°F (-10°C).

PESTICIDE DISPOSAL

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING

Dilutable Seed Treatment Products in Non-Refillable Plastic Containers

Rigid Non-refillable containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs) Non-refillable container. Do not reuse or refill this container. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781), or rinse and either recycle or dispose of the container as follows:

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, and Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSOURCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer



PROSPER®

EverGol®

Lot No. _____ Net Contents _____

Prosper® EverGol®

For use as a fungicide and insecticide seed dressing for protection against listed soilborne, seedborne, and post-emergence diseases and insect pests of listed crop plants.

ACTIVE INGREDIENTS:	
Clothianidin	22.30%
Penflufen	0.82%
Trifloxystrobin	0.55%
Metalaxyl	0.55%
OTHER INGREDIENTS:	75.78%

Contains: 2.42 lb clothianidin; 0.09 lb penflufen;	TOTAL	100.00%
0.06 lb trifloxystrobin; 0.06 lb metalaxyl per gallon		
Contains: 290 g clothianidin; 10.7 g penflufen; 7.15 g trifloxystrobin; 7.15 g metalaxyl per liter		

EPA Reg. No. 264-1121

KEEP OUT OF REACH OF CHILDREN CAUTION

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY**

Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

Please refer to booklet for additional precautionary statements and directions for use.

FIRST AID

IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing.

CLOTHIANIDIN	GROUP	4A	INSECTICIDE
PENFLUFEN	GROUP	7	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE
METALAXYL	GROUP	4	FUNGICIDE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Store in original container away from feed and food. Store in cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 105°F (40°C) or in temperatures that fall below 14°F (-10°C).

PESTICIDE DISPOSAL

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING

Dilutable Seed Treatment Products in Non-Refillable Plastic Containers

Rigid Non-refillable containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)

Non-refillable container. Do not reuse or refill this container. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781), or rinse and either recycle or dispose of the container as follows:

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, and Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167

PROSPER and EVERGOL are registered trademarks of Bayer.

©2019 Bayer CropScience

190923D 11/19

US050523157D