

FIFRA Section 2(ee) Recommendation

EVITO™ 480 SC Fungicide

EPA Reg. No. 66330-64

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U. S. Environmental Protection Agency

FOR DISTRIBUTION ONLY IN STATES WHERE EVITO™ 480 SC Fungicide (EPA Reg. No. 66330-64) FUNGICIDE IS CURRENTLY REGISTERED

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions for use, precautions and limitations on the product label attached to the container for EVITO 480 SC Fungicide. Also observe all directions, precautions and limitations on the labeling of all other products mentioned below. This recommendation must be in the possession of the user at the time of application.

For Control or Suppression of certain diseases on Tuberous and Corm Vegetables (ex. Potato) and Fruiting Vegetables (ex. Tomato).

Crop	Disease Control	Rates to Use	Application Timing
Tuberous and Corm Vegetables (Potato, Arracacha, arrowroot, artichoke (Chinese, Jerusalem), canna (edible), cassava (bitter, sweet), chayote (root), chufa, dasheen, ginger, leren, sweet potato, tanier, turmeric, and yam (bean, true))	Early Blight (<i>Alternaria solani</i>)	2.0 to 3.8 fl oz/A (0.063 to 0.12 lbs ai/A)	Begin applications preventively and continue as needed on a 7 to 10 day interval. Use higher rates when disease pressure is severe.
	Disease Suppression		
	Late Blight (<i>Phytophthora infestans</i>)	2.0 to 3.8 fl oz/A (0.063 to 0.12 lbs ai/A)	Begin applications preventively and continue as needed on a 7 day interval. Use higher rates when disease pressure is severe. Tank-mix or alternate with a protectant fungicide at low recommended rate for late blight control.
Crop	Disease Control	Rates to Use	Application Timing
Fruiting Vegetables (Tomato, Eggplant, groundcherry (<i>Physalis</i> sp.), pepino, pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), and tomatillo)	Early Blight (<i>Alternaria solani</i>)	2.0 to 5.7 fl oz/A (0.063 to 0.18 lbs ai/A)	Begin applications preventively and continue as needed on a 7 to 10 day interval. Use higher rates when disease pressure is severe.
	Target spot (<i>Corynespora cassiicola</i>)		
	Disease Suppression		
	Late Blight (<i>Phytophthora infestans</i>)	2.0 to 5.7 fl oz/A (0.063 to 0.18 lbs ai/A)	Begin applications preventively and continue as needed on a 7 day interval. Use higher rates when disease pressure is severe. Tank-mix or alternate with a protectant fungicide at low recommended rate for late blight control.
Notes: <ul style="list-style-type: none"> Do not apply more than 22.8 fl oz (0.72 lbs ai) of EVITO 480 SC Fungicide per crop cycle. Alternate every application of a QoI fungicide with at least one application of another effective mode of action fungicide. Use of a surfactant may increase coverage and efficacy of this product. 			